

Leader election using random walks

Gerold Alsmeyer¹, Zakhar Kabluchko¹ and Alexander Marynych^{1,2}

Abstract In the classical leader election procedure all players toss coins independently and those who get tails leave the game, while those who get heads move to the next round where the procedure is repeated. We investigate a generalization of this procedure in which the labels (positions) of the players who remain in the game are determined using an integer-valued random walk. We study the asymptotics of some relevant quantities for this model such as: the positions of the persons who remained after n rounds; the total number of rounds until all the persons among $1, 2, \dots, M$ leave the game; and the number of players among $1, 2, \dots, M$ who survived the first n rounds. Our results lead to some interesting connection with Galton-Watson branching processes and with the solutions of certain stochastic-fixed point equations arising in the context of the stability of point processes under thinning. We describe the set of solutions to these equations and thus provide a characterization of one-dimensional point processes that are stable with respect to thinning by integer-valued random walks.

AMS 2000 subject classifications: primary 60F05, 60G55; secondary 60J10

Keywords: Galton-Watson branching process, leader-election procedure, random sieve, restricted self-similarity, stable point process, stochastic-fixed point equation

1 Introduction

The classical leader-election procedure [5, 11, 13, 15, 19], when applied to the infinite set of positive integers \mathbb{N} , may be viewed as a random sieve which in each round eliminates an integer not yet sieved in accordance with the outcome of a coin tossing event. The integers are typically viewed as players in a game who independently toss a coin so as to determine whether they will stay in the game for the next round or not. An alternative

¹ Inst. Math. Statistics, Department of Mathematics and Computer Science, University of Münster, Orleans-Ring 10, D-48149 Münster, Germany.

² Faculty of Cybernetics, Taras Shevchenko National University of Kyiv, 01601 Kyiv, Ukraine
e-mail: gerolda@math.uni-muenster.de, zakhar.kabluchko@uni-muenster.de, marynych@unicyb.kiev.ua




Fig. 1 A realization of the leader election procedure. The players are arranged horizontally, the vertical axis corresponds to the number of rounds, the bottom line being the first round. Players leaving (staying in) the game are shown as empty (filled) circles.

description is the following: Relabel kept integers (players) at the beginning of each round by $1, 2, \dots$ while keeping the original order. Then let $R = \{R(k) : k \geq 1\}$ be the random set of integers kept for the next round, where

$$R(0) := 0, \quad R(k) := \xi_1 + \dots + \xi_k, \quad k \geq 1$$

is a random walk with independent identically distributed (iid) increments ξ_1, ξ_2, \dots having a geometric distribution on \mathbb{N} . Adopting this viewpoint, a natural generalization is to replace the geometric distribution by an arbitrary distribution $(p_n)_{n \geq 1}$ on \mathbb{N} with $p_1 < 1$.

The purpose of this paper is to study the asymptotics of some relevant quantities for this generalization which will lead us to some interesting connection with Galton-Watson branching processes (GWP) and the solutions of certain related stochastic-fixed point equations (SFPE). Such SFPE's in turn arise in connection with the stability of point processes as will be explained in Section 2.3.

A more formal model description is next; see Figure 1 for a sample realization. Let $R^{(n)}$, $n \geq 1$, be independent copies of a random walk R and denote the increments of $R^{(n)}$ by $\xi_1^{(n)}, \xi_2^{(n)}, \dots$. That is, the $\xi_k^{(n)}$ for $k, n \in \mathbb{N}$ are iid random variables with $\mathbb{P}\{\xi_k^{(n)} = i\} = p_i$, $i \in \mathbb{N}$, and

$$R^{(n)}(0) = 0, \quad R^{(n)}(k) = \xi_1^{(n)} + \dots + \xi_k^{(n)}.$$

In round n , players with current labels $R^{(n)}(1), R^{(n)}(2), \dots$ stay for the next round while all other players leave the game. Remaining players are relabeled by $1, 2, \dots$ and the procedure is repeated over and over again. The quantities to be studied hereafter are

- $N_M^{(n)}$, the number of players among $1, 2, \dots, M$ who survived the first n rounds, formally

$$N_M^{(0)} := M \quad \text{and} \quad N_M^{(n)} := \#\{j \in \mathbb{N} : R^{(n)}(j) \leq N_M^{(n-1)}\} \quad (1)$$

for $M \in \mathbb{N}_0 := \{0, 1, 2, \dots\}$ and $n \in \mathbb{N}$.

- $1 \leq S_1^{(n)} < S_2^{(n)} < S_3^{(n)} < \dots$, the original numbers of the players who survived the first n rounds, formally

$$S_j^{(n)} := \inf\{i \in \mathbb{N} : N_i^{(n)} = j\} \quad (2)$$

for $j \in \mathbb{N}$ and $n \in \mathbb{N}_0$.

- $T(M)$, the number of rounds until all players $1, 2, \dots, M$ have been eliminated, thus

$$T(M) := \inf\{n \in \mathbb{N} : N_M^{(n)} = 0\} \quad (3)$$

for $M \in \mathbb{N}$.

The connection with simple GWP's stems from the basic observation that, for each $n \in \mathbb{N}_0$,

$$(S_1^{(n)}, S_2^{(n)}, \dots) = (S_{R^{(n)}(1)}^{(n-1)}, S_{R^{(n)}(2)}^{(n-1)}, \dots),$$

whence, using the initial conditions $S_j^{(0)} = j$ for $j \in \mathbb{N}$,

$$(S_1^{(n)}, S_2^{(n)}, \dots) = (R^{(1)} \circ \dots \circ R^{(n)}(1), R^{(1)} \circ \dots \circ R^{(n)}(2), \dots)$$

for $n \in \mathbb{N}_0$, where \circ denotes the usual composition $f \circ g(\cdot) = f(g(\cdot))$. This shows that the random vector $(S_1^{(n)}, S_2^{(n)}, \dots)$ is the n -fold forward iteration of the random walk R when viewed as a random mapping from \mathbb{N} to \mathbb{N} . Passing to the backward iterations, which does not change the distribution, we obtain the basic relation for our leader-election procedure:

$$(S_1^{(n)}, S_2^{(n)}, \dots) \stackrel{d}{=} (R^{(n)} \circ \dots \circ R^{(1)}(1), R^{(n)} \circ \dots \circ R^{(1)}(2), \dots) \quad (4)$$

for $n \in \mathbb{N}_0$. Now, the j -th coordinate of the random vector on the right-hand side is nothing but the number of descendants of individuals $1, \dots, j$ in generation n of a GWP starting from countably many individuals $1, 2, \dots$ and having offspring distribution $(p_n)_{n \geq 1}$. A sample realization of this GWP is shown in Figure 2. Since we assume $p_1 < 1$, the GWP is supercritical and survives with probability one. Two classes of leader-election procedures will be investigated separately hereafter and lead to quite different asymptotics: those generated by a law $(p_n)_{n \geq 1}$ with finite-mean, in which case the corresponding GWP has also finite mean, and those where $\sum_{n \geq 1} np_n = \infty$.

As indicated by (4), our limit results for $(S_1^{(n)}, S_2^{(n)}, \dots)$, stated as Theorems 2.1 and 2.8, will be derived from appropriate limit results for GWP's. The asymptotic behavior of $N_M^{(n)}$, as $n, M \rightarrow \infty$, and $T(M)$, as $M \rightarrow \infty$, is then found in a straightforward manner by drawing on the simple duality relations

$$\{N_M^{(n)} \geq k\} = \{S_k^{(n)} \leq M\} \quad \text{and} \quad \{T(M) \leq k\} = \{S_1^{(k)} > M\}$$

for $k, M \in \mathbb{N}$ and $n \in \mathbb{N}_0$, see Theorems 2.2, 2.3, 2.9 and 2.11. The limit processes appearing in Theorems 2.1 and 2.8 are solutions to certain stochastic fixed-point equations, see (6) and (14). The description of the set of all solutions to these equations, to be given in Theorems 2.6 and 2.13, is a much more delicate question, for the necessary analysis heavily relies on deep results about the behavior of GWP.

We have organized this work as follows. All results are stated in the next section. Proofs are then provided in Section 3 for the case when $\sum_{n \geq 1} np_n < \infty$ and in Section 4 for the case when $\sum_{n \geq 1} np_n = \infty$. Some technical lemmata may be found in a short appendix. Let us finally mention that in [1], we have studied another modification of the classical leader-election procedure which is based on records in an iid sample. This modification is closely related to the Poisson-Dirichlet coalescent and its asymptotic behavior is different from the model studied here.




Fig. 2 A Galton-Watson process starting with countably many individuals $1, 2, \dots$. The $(n+1)$ -st row (counting from the top) shows the individuals in the n -th generation. The j -th individual in generation n is shown at horizontal position $\mu^{-n}j$, where $\mu \in (1, \infty)$ is the expected number of offspring of one individual. Any individual in the n -th generation is connected by a line segment to its last descendant in generation $n+1$. For example, individual 1 in the top row has 2 direct descendants, whereas individuals $2, \dots, 8$ in the top row have one direct descendant each.

2 Results

In the following, ξ always denotes a generic copy of the increments of the random walks $R^{(n)}$ underlying the considered leader-election procedure, thus $\mathbb{P}\{\xi = n\} = p_n$ for $n \in \mathbb{N}$. Moreover, $\xrightarrow{f.d.d.}$ shall denote convergence of finite dimensional distributions.

2.1 The case when $\mathbb{E}\xi < \infty$

We start with a convergence result for $(S_j^{(n)})_{j \geq 1}$. Put $\mu := \mathbb{E}\xi$ and let $(Z_n)_{n \geq 0}$ denote a GWP with offspring distribution $(p_n)_{n \in \mathbb{N}}$. If μ is finite, thus $\mu \in (1, \infty)$, then the normalization $(\mu^{-n}Z_n)_{n \geq 0}$ constitutes a positive martingale and converges a.s. to a limit Z_∞ which is either a.s. positive, namely if $\mathbb{E}\xi \log \xi < \infty$, or zero.

Theorem 2.1 *Suppose that $\mu \in (1, \infty)$. Then, as $n \rightarrow \infty$,*

$$\left(\frac{S_1^{(n)}}{\mu^n}, \frac{S_2^{(n)}}{\mu^n}, \frac{S_3^{(n)}}{\mu^n}, \dots \right) \xrightarrow{f.d.d.} (Z_\infty^{(1)}, Z_\infty^{(1)} + Z_\infty^{(2)}, Z_\infty^{(1)} + Z_\infty^{(2)} + Z_\infty^{(3)}, \dots), \quad (5)$$

where $Z_\infty^{(1)}, Z_\infty^{(2)}, \dots$ are independent copies of Z_∞ . The distribution of the limit vector in (5) satisfies the SFPE

$$\left(\mu X_1, \mu X_2, \dots\right) \stackrel{d}{=} \left(X_{R(1)}, X_{R(2)}, \dots\right), \quad (6)$$

where the random walk $(R(j))_{j \geq 0}$ on the right-hand side with generic increment ξ is independent of $(X_j)_{j \geq 0}$.

In what follows we denote by $D := D[0, \infty)$ the Skorokhod space of real-valued functions that are defined and right-continuous on $[0, \infty)$, and with finite limits from the left on $(0, \infty)$. Weak convergence on the space D (which may be endowed with the J_1 - or M_1 -topology depending on the situation) is denoted by \xrightarrow{d} .

Theorem 2.2 *Let $\mu \in (1, \infty)$ and $\mathbb{E}\xi \log \xi < \infty$. Then, as $n \rightarrow \infty$,*

$$N_{\lfloor \mu^n \cdot \rfloor}^{(n)} \xrightarrow{d} N'(\cdot)$$

weakly in the Skorokhod space D endowed with the J_1 -topology, where $N'(x) := \#\{k \in \mathbb{N} : Z_\infty^{(1)} + \dots + Z_\infty^{(k)} \leq x\}$ for $x \geq 0$.

Note that $N'(\cdot)$ is the renewal counting process associated with $(\sum_{j=1}^k Z_\infty^{(j)})_{k \geq 1}$ and therefore a homogeneous Poisson process if the law of Z_∞ is exponential, see Example 2.5 below.

The next theorem provides a one-dimensional result for the number $T(M)$ of rounds until all players $1, \dots, M$ have been eliminated. Although not difficult to prove, we have refrained from a statement of a corresponding functional limit theorem like Theorem 2.2 because it would have required the introduction of a lot more additional notation. See [1] for results of this type in a similar setup.

Theorem 2.3 *Let $\mu \in (1, \infty)$ and $\mathbb{E}\xi \log \xi < \infty$. For fixed $x > 0$, we have*

$$T(\lfloor \mu^n x \rfloor) - n \xrightarrow{d} T'(x),$$

where the distribution of $T'(x)$ is given by

$$\mathbb{P}\{T'(x) \leq k\} = \mathbb{P}\{Z_\infty > \mu^{-k} x\}, \quad k \in \mathbb{Z}. \quad (7)$$

Remark 2.4 In Theorems 2.2 and 2.3, we assume $\mathbb{E}\xi \log \xi < \infty$ since otherwise $Z_\infty = 0$ a.s. which means that $N'(x)$ and $T'(x)$ are undefined. If $\mu \in (1, \infty)$ and $\mathbb{E}\xi \log \xi = \infty$ it is possible to obtain counterparts of Theorems 2.1, 2.2 and 2.3 using a Seneta-Heyde normalization for the GWP $(Z_n)_{n \geq 0}$. More precisely, if $(c_n)_{n \in \mathbb{N}_0}$ is such that, as $n \rightarrow \infty$,

$$\frac{Z_n}{c_n} \rightarrow Z_\infty^{(SH)} \quad \text{a.s.}$$

where $Z_\infty^{(SH)}$ is a.s. positive, then all the claims of Theorems 2.1, 2.2 and 2.3 remain valid upon dropping the assumption $\mathbb{E}\xi \log \xi < \infty$ and after replacing μ^n by c_n and Z_∞ by $Z_\infty^{(SH)}$ everywhere, except the term μ^{-k} in formula (7). The latter formula in that case takes the form

$$\mathbb{P}\{T'(x) \leq k\} = \mathbb{P}\{Z_\infty^{(SH)} > \mu^{-k}x\}, \quad k \in \mathbb{Z}.$$

Example 2.5 In the classical leader-election procedure, the players who stay in the game for the next round are determined by iid Bernoulli trials, so that ξ has a geometric distribution on \mathbb{N} with some parameter $p \in (0, 1)$. Looking at the number of players who survive the first n rounds then leads to a Bernoulli process with parameter p^n . It follows that the process on the right-hand side of (5) is a random walk with standard exponential increments $Z_\infty^{(k)}$, $k \in \mathbb{N}$, and $(N'(x))_{x \geq 0}$ a standard Poisson process.

An interesting and intriguing problem arising from Theorem 2.1 is to describe the set of all positive nondecreasing solutions to the SFPE (6). Indeed, we can view (6) as a definition of a certain stability property of point processes; see Section 2.3 for more details. Note that if the distribution of a random sequence (X_1, X_2, \dots) satisfies (6) and $G : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, where $\mathbb{R}^+ := [0, \infty)$, is an arbitrary nondecreasing random process independent of (X_1, X_2, \dots) and with the restricted self-similarity property

$$(G(\mu t))_{t \in \mathbb{R}^+} \stackrel{f.d.d.}{=} (\mu G(t))_{t \in \mathbb{R}^+}, \quad (8)$$

then $(G(X_1), G(X_2), \dots)$ also satisfies (6). The class of such processes forms a semigroup with respect to composition of independent realizations. Our next result shows that all solutions to (6) can be constructed in the above way and the limit in (5) provides a solution which is fundamental in a certain sense.

Theorem 2.6 *Let (X_1, X_2, \dots) be a random element of $\mathbb{R}^{\mathbb{N}}$ such that $0 \leq X_1 \leq X_2 \leq \dots$ and (6) holds with $(R(j))_{j \geq 0}$ independent of $(X_j)_{j \in \mathbb{N}}$ and $\mu = \mathbb{E}R(1) = \mathbb{E}\xi$. Assume further that*

$$\mathbb{E}R(1) \log R(1) = \mathbb{E}\xi \log \xi < \infty, \quad (9)$$

so that $\mathbb{P}(Z_\infty > 0) = 1$. Let $(Z_\infty^{(j)})_{j \in \mathbb{N}}$ be independent copies of Z_∞ . Then there exists a nondecreasing random process $(G(t))_{t \in \mathbb{R}^+}$, independent of $(Z_\infty^{(j)})_{j \geq 1}$ and satisfying (8), such that

$$(X_j)_{j \geq 1} \stackrel{d}{=} (G(Z_\infty^{(1)} + \dots + Z_\infty^{(j)}))_{j \geq 1}.$$

Remark 2.7 Our proof of Theorem 2.6 does not work in the case when $\mu \in (1, \infty)$ and $\mathbb{E}\xi \log \xi = \infty$, and we do not know whether the result still holds without the $(\xi \log \xi)$ -assumption after replacing Z_∞ by $Z_\infty^{(SH)}$ as in Remark 2.4. Of course, the direct part of the claim, namely that all vectors of the form

$$(G(Z_\infty^{(SH)(1)} + \dots + Z_\infty^{(SH)(j)}))_{j \geq 1}$$

are solutions to (6), is obvious. However, it remains open whether the converse is true.

2.2 The case when $\mathbb{E}\xi = \infty$

The behavior of GWP's with infinite mean has been studied by various authors including Darling [7], Seneta [21], Davies [8], Grey [12], Schuh and Barbour [20]. In the last reference, it has been proved that for an arbitrary infinite-mean GWP there always exists a function

U and a sequence of deterministic constants $(C_n)_{n \in \mathbb{N}}$ such that $U(Z_n)/C_n$ converges almost surely to a non-degenerate random variable. However, the implicit construction of U and C_n makes it difficult to deduce any quantitative result for our model in this general framework. Here we work under the assumption of Davies [8], which to the best of our knowledge, are the most general conditions allowing the explicit construction of U and C_n . They also guarantee that the a.s. limit is positive on the set of survival. Davies' assumption is:

$$x^{-\alpha-\gamma(x)} \leq \mathbb{P}\{\xi \geq x\} \leq x^{-\alpha+\gamma(x)}, \quad x \geq x_0, \quad (10)$$

for some $0 < \alpha < 1$, $x_0 \geq 0$, and a nonincreasing, non-negative function $\gamma(x)$ such that $x^{\gamma(x)}$ is nondecreasing and $\int_{x_0}^{\infty} \gamma(\exp(e^x)) dx < \infty$. For example, this assumption is satisfied if $x^\alpha \mathbb{P}\{\xi > x\}$ stays bounded away from 0 and $+\infty$ for $x \geq x_0$ (to see this, take $\gamma(x) = C/\log x$). For a GWP $(Z_n)_{n \geq 0}$ with generic offspring variable ξ satisfying the above assumptions and $Z_0 = 1$, Davies [8, Thms. 1 and 2] proved the existence of the limit

$$Z_\infty^* := \lim_{n \rightarrow \infty} \alpha^n \log(1 + Z_n) \in (0, \infty) \quad \text{a.s.} \quad (11)$$

Since $\mathbb{P}\{\xi = 0\} = 0$ in our setting, we have $1 \leq Z_n \rightarrow \infty$ a.s. and therefore the equivalence of (11) with

$$Z_\infty^* = \lim_{n \rightarrow \infty} \alpha^n \log Z_n \in (0, \infty) \quad \text{a.s.} \quad (12)$$

Moreover, Z_∞^* has a continuous distribution on $(0, \infty)$, see [22, p. 715] or [2, bottom of p. 3763].

Theorem 2.8 *Consider a leader-election procedure in which the distribution of ξ satisfies (10) for some $\alpha \in (0, 1)$. Then*

$$\left(\alpha^n \log S_j^{(n)} \right)_{j \geq 1} \xrightarrow{f.d.d.} \left(Z_\infty^{(*,1)}, Z_\infty^{(*,1)} \vee Z_\infty^{(*,2)}, Z_\infty^{(*,1)} \vee Z_\infty^{(*,2)} \vee Z_\infty^{(*,3)}, \dots \right), \quad (13)$$

where $Z_\infty^{(*,1)}, Z_\infty^{(*,2)}, \dots$ are independent copies of Z_∞^* and \vee denotes the maximum. The distribution of the limit vector in (13) satisfies the SFPE

$$(X_1, X_2, \dots) \stackrel{d}{=} (\alpha X_{R(1)}, \alpha X_{R(2)}, \dots), \quad (14)$$

where the random walk $(R(j))_{j \geq 1}$ on the right-hand side is independent of $(X_j)_{j \geq 1}$.

On the right-hand side of (13), we thus have the running maximum process of iid positive random variables instead of sums as in the finite-mean case. For a study of this process (including, for example, a proof of the Markov property and exact expressions for the transition probabilities), we refer to [17, Lectures 14, 15, 17]. Let us stress that this process has multiple elements which means that the original numbers of players remaining after n rounds tend to build clusters (at least asymptotically on the log-scale). In fact, it follows from the Rényi theorem on records [17, p. 58] that among the first k elements of this process there are just $\simeq \log k$ distinct ones, as $k \rightarrow \infty$.

Theorem 2.9 *Under the assumptions of Theorem 2.8, we have*

$$N_{\lfloor \exp(\cdot \alpha^{-n}) \rfloor}^{(n)} \xrightarrow{d} N''(\cdot),$$

weakly in the Skorokhod space D endowed with the M_1 -topology, where $N''(x) := \#\{k \in \mathbb{N} : Z_\infty^{(*,1)} \vee \dots \vee Z_\infty^{(*,k)} \leq x\}$, $x \geq 0$.

Remark 2.10 The above theorem breaks down if D is endowed with the J_1 -topology. Indeed, the sample paths of the process $(N_{\lfloor \exp(x\alpha^{-n}) \rfloor}^{(n)})_{x \geq 0}$ belong to the set of piecewise constant non-decreasing functions with jumps of size 1 which is a closed subset of D in the J_1 -topology. But the process N'' has jumps of size at least 2 with probability 1 (due to the clustering), hence Theorem 2.9 cannot hold when using the J_1 -topology.

The next result is the counterpart of Theorem 2.3 in the infinite-mean case.

Theorem 2.11 *Under the assumptions of Theorem 2.8, we have, for any fixed $x > 0$,*

$$T([e^{\alpha^{-n}x}]) - n \xrightarrow{d} T''(x)$$

as $n \rightarrow \infty$, where the distribution of $T''(x)$ is given by

$$\mathbb{P}\{T''(x) \leq k\} = \mathbb{P}\{Z_\infty^* > \alpha^k x\}, \quad k \in \mathbb{Z}.$$

An interesting example of an infinite-mean GWP is obtained by choosing the law of ξ , i.e. $(p_n)_{n \geq 1}$, to be a Sibuya distribution with generating function

$$f_\alpha(t) := \mathbb{E}t^\xi = 1 - (1-t)^\alpha, \quad 0 \leq t \leq 1,$$

for some parameter $\alpha \in (0, 1)$.

Proposition 2.12 *If ξ has a Sibuya distribution with parameter $\alpha \in (0, 1)$, then weak convergence of point processes on \mathbb{R}^+ holds true, viz.*

$$\sum_{j=1}^{\infty} \delta_{\alpha^n \log S_j^{(n)}} \xrightarrow{w} \sum_{i=1}^{\infty} G_i \delta_{P_i},$$

where $P_1 < P_2 < \dots$ are the points of a standard Poisson process on $(0, \infty)$, and, given these points, the random variables G_1, G_2, \dots are conditionally independent with G_j having a geometric distribution with parameter e^{-P_j} .

So we have in the Sibuya case that after normalization with $\alpha^n \log x$ and for large n , the points $S_j^{(n)}$ form approximately a standard Poisson process and have geometrically distributed multiplicities (cluster sizes), their parameters being $\approx (S_j^{(n)})^{-\alpha^n}$.

Our last theorem is the infinite-mean counterpart of Theorem 2.6 and provides the description of the set of all solutions to (14) under the Davies' assumption (10) on ξ .

Theorem 2.13 *Let (X_1, X_2, \dots) be a random element of $\mathbb{R}^{\mathbb{N}}$ such that $0 \leq X_1 \leq X_2 \leq \dots$ and (14) holds with $(R(j))_{j \geq 0}$ independent of $(X_j)_{j \in \mathbb{N}}$. Suppose further Davies' condition (10) and let $(Z_\infty^{(*,j)})_{j \in \mathbb{N}}$ be independent copies of Z_∞^* , defined by (12). Then there exists a nondecreasing random process $G : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ satisfying*

$$(G(\alpha t))_{t \in \mathbb{R}^+} \stackrel{d}{=} (\alpha G(t))_{t \in \mathbb{R}^+} \tag{15}$$

and independent of $(Z_\infty^{(*,j)})_{j \geq 1}$ such that

$$(X_j)_{j \in \mathbb{N}} \stackrel{d}{=} \left(G(Z_\infty^{(*,1)} \vee \dots \vee Z_\infty^{(*,j)}) \right)_{j \geq 1}.$$

2.3 Stability of point processes: a natural connection

The description of the set of solutions to the fixed-point equations (6) and (14) provided by our Theorems 2.6 and 2.13 may be interpreted from a different point of view involving the notion of stability of point processes, see [9, 10] and also a recent preprint [24]. To define such stability usually requires two operations, namely *thinning* and *rescaling*. Given a point process $\mathcal{X} := \sum_{k=1}^{\infty} \delta_{X_k}$ in $[0, \infty)$ with $0 \leq X_1 \leq X_2 \leq \dots$, and an increasing integer-valued random walk $R = (R(k))_{k \geq 1}$, we define the thinning of \mathcal{X} by R as

$$\mathcal{X} \bullet R := \sum_{k=1}^{\infty} \delta_{X_{R(k)}}.$$

This random operation transforms \mathcal{X} into a “sparser” point process $\mathcal{X} \bullet R$ by removing points of \mathcal{X} with indices outside the range of the random walk R . In order to compensate such thinning, a second operation is used for rescaling, namely the usual multiplication $a \cdot \mathcal{X} := \sum_{k=1}^{\infty} \delta_{aX_k}$, $a \in (0, 1)$. We call a point process \mathcal{X} *a-stable with respect to thinning by an integer-valued increasing random walk R* if

$$\mathcal{X} \stackrel{d}{=} a \cdot (\mathcal{X} \bullet R). \tag{16}$$

Note that \mathcal{X} is *a-stable* if and only if \mathcal{X}^β is a^β -stable, where $\mathcal{X}^\beta := \sum_{k=1}^{\infty} \delta_{X_k^\beta}$ and $\beta > 0$. This observation implies that, given a random walk R , it is enough to study only *a-stable* point processes for some particular choice of $a \in (0, 1)$.

Adopting this viewpoint, Theorems 2.6 and 2.13 are nothing else but characterizations of *a-stable* point processes with respect to thinning by random walks. Moreover, the particular choice of a ($a = \mu^{-1}$ in Theorem 2.6 and $a = \alpha$ in Theorem 2.13) does not reduce generality which means that the aforementioned theorems actually provide the description of the set of solutions to (16) for arbitrary $a \in (0, 1)$.

3 Proofs in the finite-mean case

3.1 Auxiliary results about GWP’s with finite-mean offspring distribution

The results of this section are used in the proof of Theorem 2.6.

Lemma 3.1 *Let θ be a positive random variable with $\mu := \mathbb{E}\theta \in (1, \infty)$ and $\mathbb{E}\theta \log^+ \theta < \infty$. Then, for each $p \in [1, 2)$,*

$$\sum_{k=1}^{\infty} \left(\mathbb{E} |\theta - \mu| \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}} \right)^{1/p} < \infty.$$

Proof. The statement is obvious for $p = 1$ as the series on the left-hand side then reduces to $\mathbb{E} |\theta - \mu| \mathbf{1}_{\{|\theta - \mu| > 1\}}$. So let $p \in (1, 2)$ and choose $q > 2 > p$ such that $1/p + 1/q = 1$. Using Hölder's inequality $|\sum_k a_k b_k| \leq (\sum_k |a_k|^p)^{1/p} (\sum_k |b_k|^q)^{1/q}$ with $b_k = k^{-1/p}$ and $a_k = \left(k \mathbb{E} |\theta - \mu| \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}} \right)^{1/p}$, we obtain

$$\begin{aligned} & \sum_{k=1}^{\infty} \left(\mathbb{E} |\theta - \mu| \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}} \right)^{1/p} \\ & \leq \left(\sum_{k=1}^{\infty} k \mathbb{E} |\theta - \mu| \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}} \right)^{1/p} \left(\sum_{k=1}^{\infty} k^{-q/p} \right)^{1/q}. \end{aligned}$$

The second series converges because $q > p$ and the first one can be bounded by

$$\mathbb{E} \left(\sum_{k=1}^{\infty} \left(\frac{\log |\theta - \mu|}{\log \mu} + 1 \right) |\theta - \mu| \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}} \right)$$

which is finite because $\mathbb{E} \theta \log^+ \theta < \infty$. \square

Lemma 3.2 *Let $(\theta_k^{(n)})_{n \in \mathbb{N}, k \in \mathbb{N}}$ be an array of independent copies of a positive random variable θ having $\mu = \mathbb{E} \theta \in (1, \infty)$ and $\mathbb{E} \theta \log^+ \theta < \infty$. For $n \in \mathbb{N}$, define the increasing random walks*

$$R^{(n)}(0) := 0, \quad R^{(n)}(k) := \theta_1^{(n)} + \dots + \theta_k^{(n)}, \quad k \in \mathbb{N}.$$

Then, for any $T > 0$,

$$\sum_{n=1}^{\infty} \sup_{t \in [0, T]} \left| \mu^{-n} R^{(n)}(\lfloor t \mu^{n-1} \rfloor) - t \right| < \infty \quad a.s.$$

Proof. Defining $\theta_{\leq, k}^{(n)} := (\theta_k^{(n)} - \mu) \mathbf{1}_{\{\theta_k^{(n)} - \mu \leq \mu^n\}}$ and $\theta_{>, k}^{(n)} := (\theta_k^{(n)} - \mu) \mathbf{1}_{\{\theta_k^{(n)} - \mu > \mu^n\}}$, we have

$$\begin{aligned} \sup_{t \in [0, T]} \left| \mu^{-n} R^{(n)}(\lfloor t \mu^{n-1} \rfloor) - t \right| & \leq \mu^{-n} \sup_{t \in [0, T]} |R^{(n)}(\lfloor t \mu^{n-1} \rfloor) - \mu \lfloor t \mu^{n-1} \rfloor| \\ & \quad + \mu^{-n} \sup_{t \in [0, T]} |\mu \lfloor t \mu^{n-1} \rfloor - t \mu^n| \\ & \leq \mu^{-n} \left(\sup_{t \in [0, T]} \left| \sum_{k=1}^{\lfloor t \mu^{n-1} \rfloor} \theta_{\leq, k}^{(n)} \right| + \sup_{t \in [0, T]} \left| \sum_{k=1}^{\lfloor t \mu^{n-1} \rfloor} \theta_{>, k}^{(n)} \right| \right) + \mu^{-n}. \end{aligned}$$

Hence it is enough to show that

$$\sum_{n=1}^{\infty} \mu^{-n} \sup_{0 \leq m \leq \lfloor T\mu^{n-1} \rfloor} \left| \sum_{k=1}^m \theta_{\leq, k}^{(n)} \right| \quad \text{and} \quad \sum_{n=1}^{\infty} \mu^{-n} \sup_{0 \leq m \leq \lfloor T\mu^{n-1} \rfloor} \left| \sum_{k=1}^m \theta_{>, k}^{(n)} \right| \quad (17)$$

are both almost surely finite. As for the second series, this follows by the Borel-Cantelli lemma if we can show that

$$\sum_{n=1}^{\infty} \mathbb{P} \left\{ \text{there exists } k = 1, \dots, \lfloor T\mu^{n-1} \rfloor : \theta_{>, k}^{(n)} \neq 0 \right\} < \infty. \quad (18)$$

To this end, we use Boole's inequality to infer

$$\begin{aligned} \sum_{n=1}^{\infty} \mathbb{P} \left\{ \text{there exists } k = 1, \dots, \lfloor T\mu^{n-1} \rfloor : \theta_{>, k}^{(n)} \neq 0 \right\} &\leq \sum_{n=1}^{\infty} (T\mu^{n-1}) \mathbb{P} \left\{ \theta_{>, 1}^{(n)} \neq 0 \right\} \\ &= \sum_{n=1}^{\infty} (T\mu^{n-1}) \mathbb{P} \{ |\theta - \mu| > \mu^n \} = \mathbb{E} \left(\sum_{n=1}^{\infty} (T\mu^{n-1}) \mathbf{1}_{\{|\theta - \mu| > \mu^n\}} \right) \\ &\leq \frac{T}{\mu} \mathbb{E} \left(\sum_{n=1}^{\infty} |\theta - \mu| \mathbf{1}_{\{|\theta - \mu| > \mu^n\}} \right) \leq \frac{T}{\mu} \mathbb{E} \left((\theta + \mu) \sum_{n=1}^{\infty} \mathbf{1}_{\{\theta \geq \mu^n\}} \right) \\ &\leq \frac{T}{\mu} \mathbb{E} \left((\theta + \mu) \sum_{n=1}^{\infty} \mathbf{1}_{\{\log^+ \theta \geq n \log \mu\}} \right) \leq \frac{T}{\mu} \mathbb{E} \left((\theta + \mu) \frac{\log^+ \theta}{\log \mu} \right) < \infty \end{aligned}$$

where the finiteness of the last term follows from $\mathbb{E}\theta \log^+ \theta < \infty$. To show that the first series in (17) converges, we argue as follows:

$$\begin{aligned} \sum_{n=1}^{\infty} \mu^{-n} \sup_{0 \leq m \leq \lfloor T\mu^{n-1} \rfloor} \left| \sum_{k=1}^m \theta_{\leq, k}^{(n)} \right| \\ \leq \sum_{n=1}^{\infty} \mu^{-n} \sup_{0 \leq m \leq \lfloor T\mu^{n-1} \rfloor} \left| \sum_{k=1}^m \theta_{\leq, k}^{(n)} - m \mathbb{E} \theta_{\leq, 1}^{(n)} \right| + \frac{T}{\mu} \sum_{n=1}^{\infty} |\mathbb{E} \theta_{\leq, 1}^{(n)}|. \end{aligned}$$

The last term on the right-hand side is finite because

$$\sum_{n=1}^{\infty} |\mathbb{E} \theta_{\leq, 1}^{(n)}| = \sum_{n=1}^{\infty} |\mathbb{E} \theta_{>, 1}^{(n)}| \leq \sum_{n=1}^{\infty} \mathbb{E} |\theta - \mu| \mathbf{1}_{\{|\theta - \mu| > \mu^n\}} < \infty,$$

where the finiteness of the last sum has already been shown above. To bound the first term, note that $(\sum_{k=1}^m (\theta_{\leq, k}^{(n)} - \mathbb{E} \theta_{\leq, 1}^{(n)}))_{m \in \mathbb{N}}$ is an L^p -martingale for each $p \in (1, 2]$ whence

$$\begin{aligned} \mathbb{E} \left(\sup_{0 \leq m \leq \lfloor T\mu^{n-1} \rfloor} \left| \sum_{k=1}^m (\theta_{\leq, k}^{(n)} - \mathbb{E} \theta_{\leq, 1}^{(n)}) \right| \right) &\leq \left\| \sup_{0 \leq m \leq \lfloor T\mu^{n-1} \rfloor} \left| \sum_{k=1}^m (\theta_{\leq, k}^{(n)} - \mathbb{E} \theta_{\leq, 1}^{(n)}) \right| \right\|_p \\ &\leq \frac{p}{p-1} \left\| \sum_{k=1}^{\lfloor T\mu^{n-1} \rfloor} (\theta_{\leq, k}^{(n)} - \mathbb{E} \theta_{\leq, 1}^{(n)}) \right\|_p \leq \frac{2p}{p-1} T^{1/p} \mu^{(n-1)/p} \left\| \theta_{\leq, 1}^{(n)} - \mathbb{E} \theta_{\leq, 1}^{(n)} \right\|_p \end{aligned}$$

$$\leq \frac{4p}{p-1} T^{1/p} \mu^{n/p} \left\| \theta_{\leq,1}^{(n)} \right\|_p,$$

having utilized the inequalities by Doob and von Bahr-Esseen [23, Formula 4]. Put $q := \frac{p}{p-1}$, thus $1/p + 1/q = 1$, and use the inequality $|\sum_i x_i|^{1/p} \leq \sum_i |x_i|^{1/p}$ to infer

$$\begin{aligned} \sum_{n=1}^{\infty} \mu^{-n/q} \left\| \theta_{\leq,1}^{(n)} \right\|_p &\leq \sum_{n=1}^{\infty} \mu^{-n/q} \left(1 + \sum_{k=1}^n \mathbb{E} |\theta - \mu|^p \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}} \right)^{1/p} \\ &\leq \sum_{n=1}^{\infty} \mu^{-n/q} \left(1 + \sum_{k=1}^n \mu^{k/q} (\mathbb{E} |\theta - \mu| \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}})^{1/p} \right) \end{aligned}$$

and then further (with C denoting a suitable finite positive constant)

$$\begin{aligned} &\sum_{n=1}^{\infty} \mu^{-n/q} \sum_{k=1}^n \mu^{k/q} (\mathbb{E} |\theta - \mu| \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}})^{1/p} \\ &= \sum_{k=1}^{\infty} \mu^{k/q} (\mathbb{E} |\theta - \mu| \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}})^{1/p} \sum_{n=k}^{\infty} \mu^{-n/q} \\ &\leq C \sum_{k=1}^{\infty} (\mathbb{E} |\theta - \mu| \mathbf{1}_{\{\mu^{k-1} < |\theta - \mu| \leq \mu^k\}})^{1/p}, \end{aligned}$$

which is finite by Lemma 3.1 if $p \in (1, 2)$. We thus arrive at

$$\mathbb{E} \sum_{n=1}^{\infty} \mu^{-n} \sup_{0 \leq m \leq \lfloor T \mu^{n-1} \rfloor} \left| \sum_{k=1}^m \theta_{\leq,k}^{(n)} - m \mathbb{E} \theta_{\leq,1}^{(n)} \right| < \infty$$

and this completes the proof of the lemma. \square

Lemma 3.2 allows us to prove the following proposition which is the key ingredient to the proof of Theorem 2.6.

Proposition 3.3 *Let $(R^{(n)}(k))_{k \in \mathbb{N}_0, n \in \mathbb{N}}$ be as in Lemma 3.2 with θ taking positive integer values only and put*

$$g_n(t) := \mu^{-n} R^{(n)}(\lfloor \mu^{n-1} t \rfloor) \quad (19)$$

for $n \in \mathbb{N}$ and $t \geq 0$.

(A1) *There exists a D -valued random process $(Z_{\infty}(t))_{t \geq 0}$ such that*

$$(g_n \circ \dots \circ g_1(t))_{t \geq 0} \xrightarrow{n \rightarrow \infty} (Z_{\infty}(t))_{t \geq 0} \quad \text{a.s.}$$

in the space D endowed with the J_1 -topology. The process $(Z_{\infty}(t))_{t \geq 0}$ is the limit of the normalized number of descendants of individuals $1, \dots, \lfloor t \rfloor$ in a GWP with generic offspring variable θ and countably many ancestors $1, 2, \dots$, thus

$$Z_{\infty}(t) = Z_{\infty}^{(1)} + \dots + Z_{\infty}^{(\lfloor t \rfloor)}, \quad t \geq 0,$$

where the $Z_\infty^{(j)}$, $j \in \mathbb{N}$, denote independent copies of Z_∞ , the limit of the same normalized GWP with one ancestor.

(A2) For each $k \in \mathbb{N}_0$, there exists a copy $(Z_{k,\infty}(t))_{t \geq 0}$ of $(Z_\infty(t))_{t \geq 0}$ such that

$$(g_n \circ \cdots \circ g_{k+1}(t))_{t \geq 0} \xrightarrow{n \rightarrow \infty} (\mu^{-k} Z_{k,\infty}(t\mu^k))_{t \geq 0} \quad \text{a.s.}$$

in the space D endowed with the J_1 -topology.

(A3) As $t \rightarrow \infty$,

$$\inf_{k \geq 1} g_k \circ \cdots \circ g_1(t) \rightarrow \infty \quad \text{a.s.}$$

(A4) For every fixed $T > 0$,

$$\sup_{k \geq 1} \sup_{t \in [0, T]} |g_{n+k} \circ \cdots \circ g_{n+1}(t) - t| \xrightarrow{n \rightarrow \infty} 0 \quad \text{a.s.}$$

(A5) The set $\{\mu^{-k} Z_{k,\infty}(t\mu^k) : t \geq 0, k \in \mathbb{N}\}$ is almost surely dense in $[0, \infty)$.

Remark 3.4 To motivate the definition of g_n in (19), consider a GWP starting from countably many individuals $1, 2, \dots$, as shown in Figure 2. Since the expected number of offspring of one individual is μ , it is natural to place the j -th individual in generation n at position $\mu^{-n}j$. By definition of g_n , we have $g_n(\mu^{-(n-1)}j) = \mu^{-n}R^{(n)}(j)$ for all $j \in \mathbb{N}$, so that g_n maps the position of the j -th individual in generation $n-1$ to the position of its last offspring in generation n . By the law of large numbers, $g_n(t)$ should be close to t for large n . Part (A4) of Proposition 3.3 provides the confirmation of this in a very strong uniform sense, and it may also be assessed graphically in Figure 2 where slope line segments connecting individuals in consecutive generations become closer and closer to vertical lines as n grows.

Proof (of Proposition 3.3). (A1) Since $g_n \circ \cdots \circ g_1(t) = \mu^{-n}R^{(n)} \circ \cdots \circ R^{(1)}(\lfloor t \rfloor)$, the assertion is equivalent to

$$\left(\mu^{-n}R^{(n)} \circ \cdots \circ R^{(1)}(j) \right)_{j \in \mathbb{N}_0} \xrightarrow{n \rightarrow \infty} (Z_\infty(j))_{j \in \mathbb{N}_0} \quad \text{a.s.} \quad (20)$$

For each n , the sequence on the left-hand side constitutes a random walk (that is, it has iid increments) since a composition of independent increasing, \mathbb{N} -valued random walks is again a random walk (a similar statement in the theory of Lévy processes is well-known and called Bochner's subordination). Hence, in order to prove (20), it suffices to show that

$$\mu^{-n}R^{(n)} \circ \cdots \circ R^{(1)}(1) \xrightarrow{n \rightarrow \infty} Z_\infty^{(1)} \quad \text{a.s.} \quad (21)$$

The quantity on the left is the normalized number of individuals at time n in a GWP with generic offspring variable θ and one ancestor, and (21) follows from the a.s. convergence of its normalization (which is a nonnegative martingale).

(A2) This follows from (A1), when using the representation

$$g_n \circ \cdots \circ g_{k+1}(t) = \mu^{-k} \mu^{-(n-k)} R^{(n)} \circ \cdots \circ R^{(k+1)}(\lfloor t\mu^k \rfloor), \quad t \geq 0,$$

valid for $n > k$.

(A3) Here we have

$$g_k \circ \cdots \circ g_1(t) = \sum_{j=1}^{\lfloor t \rfloor} \mu^{-k} Z_k^{(j)}, \quad t \geq 0,$$

where the $(Z_k^{(j)})_{k \geq 0, j \in \mathbb{N}}$ are independent copies of a GWP $(Z_k)_{k \geq 0}$ with generic offspring variable θ and $Z_0 = 1$. Passing to the infimum yields

$$\inf_{k \in \mathbb{N}} g_k \circ \cdots \circ g_1(t) \geq \sum_{j=1}^{\lfloor t \rfloor} \inf_{k \geq 1} (\mu^{-k} Z_k^{(j)}), \quad t \geq 0$$

and the result follows from $\mathbb{P}\{\inf_{k \geq 1} (\mu^{-k} Z_k^{(1)}) = 0\} = 0$.

(A4) Fix $T > 0$. From (A3), we know that there exists a random $T_1 > 0$ such that

$$\inf_{k \geq 1} g_k \circ \cdots \circ g_1(T_1) \geq T \quad \text{a.s.}$$

We further note that

$$\begin{aligned} A_1 &:= \sup_{k \geq 0} \sup_{n > k} \sup_{t \in [0, T]} g_n \circ \cdots \circ g_{k+1}(t) = \sup_{k \geq 0} \sup_{n > k} g_n \circ \cdots \circ g_{k+1}(T) \\ &\leq \sup_{k \geq 0} \sup_{n > k} g_n \circ \cdots \circ g_1(T_1) = \sup_{n \geq 1} g_n \circ \cdots \circ g_1(T_1) < \infty \quad \text{a.s.}, \end{aligned}$$

where the finiteness is ensured by (A1), and that

$$\begin{aligned} &|g_{n+k} \circ \cdots \circ g_{n+1}(t) - t| \\ &\leq |g_{n+1}(t) - t| + \sum_{j=2}^k |g_{n+j} \circ \cdots \circ g_{n+1}(t) - g_{n+j-1} \circ \cdots \circ g_{n+1}(t)| \end{aligned}$$

by the triangle inequality. Consequently,

$$\begin{aligned} &\sup_{k \in \mathbb{N}} \sup_{t \in [0, T]} |g_{n+k} \circ \cdots \circ g_{n+1}(t) - t| \\ &\leq \sup_{t \in [0, T]} |g_{n+1}(t) - t| \\ &\quad + \sum_{j=2}^{\infty} \sup_{t \in [0, T]} |g_{n+j} \circ \cdots \circ g_{n+1}(t) - g_{n+j-1} \circ \cdots \circ g_{n+1}(t)| \\ &\leq \sup_{t \in [0, T]} |g_{n+1}(t) - t| + \sum_{j=2}^{\infty} \sup_{s \in [0, A_1]} |g_{n+j}(s) - s| \xrightarrow{n \rightarrow \infty} 0 \end{aligned}$$

by Lemma 3.2.

(A5) By (A4), for any $s \in [0, \infty)$ and $\varepsilon > 0$, there exists a random $m \in \mathbb{N}$ such that

$$s - \varepsilon/2 \leq g_{m+k} \circ \cdots \circ g_{m+1}(s) \leq s + \varepsilon/2 \quad (22)$$

for all $k \in \mathbb{N}$. Choosing k sufficiently large and applying (A2), we obtain

$$\mu^{-m} Z_{m,\infty}(s\mu^m) - \varepsilon/2 \leq g_{m+k} \circ \dots \circ g_{m+1}(s) \leq \mu^{-m} Z_{m,\infty}(s\mu^m) + \varepsilon/2. \quad (23)$$

Finally, (22) and (23) imply

$$s - \varepsilon \leq \mu^{-m} Z_{m,\infty}(s\mu^m) \leq s + \varepsilon$$

and the proof is complete. \square

3.2 Proof of Theorems 2.1, 2.2, 2.3 and 2.6

Define a random map $\psi : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ by

$$\psi((x_1, x_2, \dots)) := \frac{1}{\mu} (x_{R(1)}, x_{R(2)}, \dots) \quad (24)$$

and further $\psi_n : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ for $n \in \mathbb{N}$ by

$$\psi_n((x_1, x_2, \dots)) := \frac{1}{\mu} (x_{R^{(n)}(1)}, x_{R^{(n)}(2)}, \dots) \quad (25)$$

which are independent copies of ψ .

Proof (of Theorem 2.1). In view of the basic identity (4), we have

$$\begin{aligned} \left(\frac{S_1^{(n)}}{\mu^n}, \frac{S_2^{(n)}}{\mu^n}, \dots \right) &\stackrel{d}{=} \left(\mu^{-n} R^{(n)} \circ \dots \circ R^{(1)}(1), \mu^{-n} R^{(n)} \circ \dots \circ R^{(1)}(2), \dots \right) \\ &= \psi^{(1)} \circ \dots \circ \psi^{(n)}(1, 2, \dots). \end{aligned}$$

By (A1) of Proposition 3.3, the last sequence converges a.s. to $(Z_\infty^{(1)}, Z_\infty^{(1)} + Z_\infty^{(2)}, \dots)$ which proves (5). The fixed-point relation (6) follows from the almost sure continuity of the map ψ with respect to the product topology on $\mathbb{R}^{\mathbb{N}}$ and the continuous mapping theorem. This completes the proof of Theorem 2.1. \square

Proof (of Theorem 2.2). Observe that $(N_{\lfloor x\mu^n \rfloor}^{(n)})_{x \geq 0}$ is the counting process associated with the increasing random sequence $(\mu^{-n} S_k^{(n)})_{k \in \mathbb{N}}$. By Theorem 2.1, the finite-dimensional distributions of the latter sequence converge as $n \rightarrow \infty$ to those of the strictly increasing random sequence $(Z_\infty^{(1)} + \dots + Z_\infty^{(k)})_{k \in \mathbb{N}}$ with associated counting process $(N'(x))_{x \geq 0}$. Also, for every fixed $n \in \mathbb{N}$,

$$\lim_{k \rightarrow \infty} S_k^{(n)} = \lim_{k \rightarrow \infty} (Z_\infty^{(1)} + \dots + Z_\infty^{(k)}) = +\infty \quad \text{a.s.}$$

because $\xi > 0$ a.s. and $Z_\infty^{(1)} > 0$ a.s. By Lemma 5.2 from the Appendix, this implies the weak convergence of the corresponding counting processes on the Skorokhod space D endowed with the J_1 -topology.

Proof (of Theorem 2.3). For this result, it suffices to note

$$\begin{aligned} \mathbb{P}\{T(\lfloor \mu^n x \rfloor) - n \leq k\} &= \mathbb{P}\left\{S_1^{(n+k)} > \lfloor \mu^n x \rfloor\right\} \\ &= \mathbb{P}\left\{S_1^{(n+k)} > \mu^n x\right\} = \mathbb{P}\left\{\mu^{-(n+k)} S_1^{(n+k)} > \mu^{-k} x\right\}. \quad \square \end{aligned}$$

Proof (of Theorem 2.6). Let $(X_1^{(0)}, X_2^{(0)}, \dots)$ be a solution to (6), i.e.

$$(\mu X_1^{(0)}, \mu X_2^{(0)}, \dots) \stackrel{d}{=} (X_{R(1)}^{(0)}, X_{R(2)}^{(0)}, \dots).$$

By the Kolmogorov consistency theorem, the underlying probability space $(\Omega, \mathfrak{A}, \mathbb{P})$ may be assumed to be large enough to carry the following objects:

- the random sequence $(X_1^{(0)}, X_2^{(0)}, \dots)$;
- a two-sided sequence $(R^{(n)}(\cdot))_{n \in \mathbb{Z}}$ of independent copies of the random walk $R(\cdot)$ and the corresponding sequence of random maps (25);
- a two-sided stationary sequence $(V_k)_{k \in \mathbb{Z}} = ((X_1^{(k)}, X_2^{(k)}, \dots))_{k \in \mathbb{Z}}$ such that V_k is independent of $(\psi_n)_{n \leq k}$ for each $k \in \mathbb{Z}$, and

$$V_0 := (X_1^{(0)}, X_2^{(0)}, \dots) \quad \text{and} \quad V_k = \psi_{k+1}(V_{k+1}), \quad k \in \mathbb{Z},$$

thus

$$(X_1^{(k)}, X_2^{(k)}, \dots) = \frac{1}{\mu} (X_{R^{(k+1)}(1)}^{(k+1)}, X_{R^{(k+1)}(2)}^{(k+1)}, \dots), \quad k \in \mathbb{Z}.$$

By construction, $V_k \stackrel{d}{=} V_0$ for $k \in \mathbb{Z}$. Define a sequence of random measures $(\nu_n)_{n \geq 0}$ on $[0, \infty)$ by¹

$$\nu_n[0, t] := \begin{cases} 0, & \text{if } t < \mu^{-n}, \\ \mu^{-n} X_{\lfloor t \mu^n \rfloor}^{(n)}, & \text{if } t \geq \mu^{-n}, \end{cases}$$

which is possible because $(X_j^{(0)})_{j \geq 1}$ is nondecreasing and $(X_j^{(n)})_{j \geq 1} \stackrel{d}{=} (X_j^{(0)})_{j \geq 1}$ for each $n \in \mathbb{N}_0$. Let us assume for a moment that ν_n converges almost surely, as $n \rightarrow \infty$, to some limit random measure ν_∞ in the vague topology on $[0, \infty)$, i.e.

$$\nu_n \xrightarrow{v} \nu_\infty \quad \text{a.s.} \quad (26)$$

Let us show that $(X_k^{(0)})_{k \geq 1} \stackrel{d}{=} (G(Z_\infty^{(1)}), G(Z_\infty^{(1)} + Z_\infty^{(2)}), \dots)$ with $G(t) := \nu_\infty[0, t]$. Indeed,

$$\begin{aligned} (X_1^{(0)}, X_2^{(0)}, \dots) &= \mu^{-n} (X_{\lfloor \mu^n \widehat{Z}_n(1) \rfloor}^{(n)}, X_{\lfloor \mu^n \widehat{Z}_n(2) \rfloor}^{(n)}, \dots) \\ &= (\nu_n[0, \widehat{Z}_n(1)], \nu_n[0, \widehat{Z}_n(2)], \dots), \quad n \in \mathbb{N}, \end{aligned}$$

where $\widehat{Z}_n(j) := \mu^{-n}(R^{(n)} \circ \dots \circ R^{(1)})(j)$ is independent of ν_n . As already pointed out,

$$\left(\widehat{Z}_n(j)\right)_{j \geq 1} \xrightarrow[n \rightarrow \infty]{\mathfrak{F}} \left(Z_\infty^{(1)} + \dots + Z_\infty^{(j)}\right)_{j \geq 1} \quad \text{a.s.} \quad (27)$$

¹ For ease of notation, we write $\nu_n[0, t]$ instead of $\nu_n([0, t])$.

which in combination with (26) implies

$$\left(\nu_n, \left(\widehat{Z}_n(j) \right)_{j \in \mathbb{N}} \right) \xrightarrow{n \rightarrow \infty} \left(\nu_\infty, \left(Z_\infty^{(1)} + \dots + Z_\infty^{(j)} \right)_{j \in \mathbb{N}} \right) \quad \text{a.s.}$$

in the product topology, the components of the limit vector on the right-hand side being independent. Condition (9) entails that the law of $Z_\infty^{(1)}$ is absolutely continuous, see [3, Corollary 4 on p. 36], whence

$$\mathbb{P} \left\{ \nu_\infty(\{Z_\infty^{(1)} + \dots + Z_\infty^{(j)}\}) = 0 \right\} = 1$$

for all $j \in \mathbb{N}$. Lemma 5.1 in the Appendix now yields

$$\begin{aligned} (X_1^{(0)}, X_2^{(0)}, \dots) &= \left(\nu_n[0, \widehat{Z}_n(1)], \nu_n[0, \widehat{Z}_n(2)], \dots \right) \\ &\xrightarrow{n \rightarrow \infty} \left(\nu_\infty[0, Z_\infty^{(1)}], \nu_\infty[0, Z_\infty^{(1)} + Z_\infty^{(2)}], \dots \right) \quad \text{a.s.,} \end{aligned}$$

which shows the asserted representation of $(X_1^{(0)}, X_2^{(0)}, \dots)$ as a solution to (6).

It remains to prove (26). Recall that $g_n(t) := \mu^{-n} R^{(n)}(\lfloor t \mu^{n-1} \rfloor)$, $n \in \mathbb{N}$, $t \geq 0$, are the random maps introduced in Proposition 3.3. We have that

$$\nu_n[0, t] = \nu_{n+1}[0, g_{n+1}(t)] \quad (28)$$

for $n \in \mathbb{N}_0$ and $t \geq 0$, and this shows that ν_{n+1} differs from ν_n by a random perturbation of time. But the latter is negligible for large n by the strong law of large numbers, viz.

$$g_{n+1}(t) \xrightarrow{n \rightarrow \infty} t \quad \text{a.s.,}$$

cf. Lemma 3.2. For arbitrary fixed $k \in \mathbb{N}_0$, iteration of (28) provides us with

$$\nu_k[0, t] = \nu_{n+k}[0, g_{n+k} \circ \dots \circ g_{k+1}(t)], \quad (29)$$

in particular

$$\nu_0[0, t] = \nu_n[0, \widehat{Z}_n(\lfloor t \rfloor)] \quad (30)$$

for $n \in \mathbb{N}$ and $t \geq 0$. Since

$$\lim_{t \rightarrow \infty} \lim_{n \rightarrow \infty} \widehat{Z}_n(\lfloor t \rfloor) = \lim_{t \rightarrow \infty} Z_\infty(t) = \infty,$$

equation (30) implies that, for each $T > 0$,

$$\sup_{n \geq 0} \nu_n[0, T] < \infty \quad \text{a.s.} \quad (31)$$

Hence, $(\nu_n)_{n \geq 0}$ is a.s. relatively compact in the vague topology (see 15.7.5 in [16]). Let $(\nu_{m_n})_{n \geq 1}$, where $(m_n)_{n \in \mathbb{N}}$ is random, be an a.s. vaguely convergent subsequence and ν'_∞ its limit. From (29), we have for every fixed $k \in \mathbb{N}_0$ and $m_n > k$ that

$$\nu_k[0, t] = \nu_{m_n}[0, g_{m_n} \circ \dots \circ g_{k+1}(t)], \quad t \geq 0. \quad (32)$$

By part (A2) of Proposition 3.3

$$g_{m_n} \circ \cdots \circ g_{k+1}(t) \xrightarrow{n \rightarrow \infty} \mu^{-k} Z_{k,\infty}(t\mu^k) \quad \text{a.s.}$$

in the space D endowed with the J_1 -topology. Sending $n \rightarrow \infty$ in (32) and applying Lemma 5.1, we obtain that a.s.

$$\nu_k[0, t] = \nu'_\infty[0, \mu^{-k} Z_{k,\infty}(t\mu^k)], \quad t \geq 0 \quad (33)$$

for every fixed $k \in \mathbb{N}_0$. By part (A5) of Proposition 3.3, the random set

$$\mathcal{S} := \{\mu^{-k} Z_{k,\infty}(t\mu^k) : t \geq 0, k \in \mathbb{N}_0\}$$

is a.s. dense in $[0, \infty)$. If ν''_∞ is another subsequential limit of $(\nu_n)_{n \in \mathbb{N}_0}$, then

$$\nu''_\infty[0, t] = \nu'_\infty[0, t]$$

for all $t \in \mathcal{S}$, and therefore $\nu''_\infty = \nu'_\infty$ a.s., proving (26).

It remains to show that the random process G satisfies the restricted self-similarity property (8). But this follows immediately from

$$\nu_n[0, \mu t] = \mu^{-n} X_{\lfloor t\mu^{n+1} \rfloor}^{(n)} \stackrel{d}{=} \mu^{-n} X_{\lfloor t\mu^{n+1} \rfloor}^{(n+1)} = \mu \nu_{n+1}[0, t], \quad t \geq \mu^{-(n+1)},$$

where the equality in law is a consequence of the stationarity of $(V_k)_{k \in \mathbb{Z}}$. \square

4 Proofs in the infinite-mean case

4.1 Some auxiliary results about infinite-mean Galton-Watson processes

Lemma 4.1 *Let θ be an \mathbb{N} -valued random variable satisfying Davies' assumption, viz.*

$$x^{-\alpha-\gamma(x)} \leq \mathbb{P}\{\theta > x\} \leq x^{-\alpha+\gamma(x)}, \quad x \geq x_0,$$

for some $x_0 > 0$, $\alpha \in (0, 1)$, and a nonincreasing nonnegative function $\gamma(x)$ such that $x^{\gamma(x)}$ is nondecreasing and $\int_{x_0}^{\infty} \gamma(\exp(e^x)) dx < \infty$. Let $(\theta_n)_{n \in \mathbb{N}}$ be a sequence of independent copies of θ , and put

$$R(0) := 0, \quad R(n) := \theta_1 + \dots + \theta_n, \quad M(n) := \max_{k=1, \dots, n} \theta_k, \quad n \in \mathbb{N}.$$

Then, for each $\varepsilon > 0$ and each $\beta \in (0, \alpha)$, there exist $c, C > 0$ such that, for all $n \in \mathbb{N}$,

$$cn^{-(1/\alpha+\varepsilon)\beta\alpha^{-1}\gamma(n^{1/\alpha+\varepsilon})} \leq \mathbb{E} \left(\frac{R(n)}{n^{1/\alpha}} \right)^\beta \leq Cn^{(1/\alpha+\varepsilon)\beta\alpha^{-1}\gamma(n^{1/\alpha+\varepsilon})}, \quad (34)$$

and

$$cn^{-(1/\alpha+\varepsilon)\beta\alpha^{-1}\gamma(n^{1/\alpha+\varepsilon})} \leq \mathbb{E} \left(\frac{M(n)}{n^{1/\alpha}} \right)^{-\beta} \leq Cn^{(1/\alpha+\varepsilon)\beta\alpha^{-1}\gamma(n^{1/\alpha+\varepsilon})}. \quad (35)$$

Proof. If θ belongs to the domain of attraction of an α -stable law, $\alpha \in (0, 1)$, equivalently if $x \mapsto \mathbb{P}\{\theta > x\}$ is regularly varying at infinity with index α , then the β -th moment of $R(n)/c(n)$, where $c(\cdot)$ is such that $\lim_{n \rightarrow \infty} n \mathbb{P}\{\theta > c(n)\} = 1$, converges to the β -th moment of the limit α -stable law for all $\beta \in (0, \alpha)$. Unfortunately, Davies' condition does not imply that θ is in the domain of attraction of an α -stable law. Yet, in some sense θ can be bounded from below and above by random variables with regularly varying tails, which is the idea employed in the following argument.

We first recall, see [8, Lemma 3], that the function $x \mapsto x^{\gamma(x)}$ is slowly varying at infinity. Pick $x_1 > x_0$ so large that $x^{-\alpha+\gamma(x)} \leq 1$ for $x \geq x_1$ and let $\bar{\theta} \geq 1$ and $\underline{\theta} \geq 1$ be random variables with distributions

$$\mathbb{P}\{\bar{\theta} > x\} := \begin{cases} 1, & \text{if } 1 \leq x < x_1, \\ \sup_{y>x} y^{-\alpha+\gamma(y)}, & \text{if } x \geq x_1; \end{cases}$$

and

$$\mathbb{P}\{\underline{\theta} > x\} := \begin{cases} x_1^{-\alpha-\gamma(x_1)}, & \text{if } 1 \leq x < x_1, \\ \inf_{y>x} y^{-\alpha-\gamma(y)}, & \text{if } x \geq x_1. \end{cases}$$

From the construction it is clear that, with \leq_{st} denoting stochastic majorization,

$$\underline{\theta} \leq_{st} \theta \leq_{st} \bar{\theta}.$$

Moreover, by Theorem 1.5.3 in [4], both $x \mapsto \mathbb{P}\{\bar{\theta} > x\}$ and $x \mapsto \mathbb{P}\{\underline{\theta} > x\}$ are regularly varying at infinity of order $-\alpha$ and therefore belong to the domain of attraction of an α -stable law.

Let $(\bar{\theta}_k)_{k \geq 1}$ and $(\underline{\theta}_k)_{k \geq 1}$ be sequences of independent copies of $\bar{\theta}$ and $\underline{\theta}$, respectively, with associated zero-delayed random walks $(\bar{R}(k))_{k \geq 0}$ and $(\underline{R}(k))_{k \geq 0}$. Further, let $(\bar{c}(n))_{n \in \mathbb{N}}$ and $(\underline{c}(n))_{n \in \mathbb{N}}$ be such that

$$\lim_{n \rightarrow \infty} n \mathbb{P}\{\underline{\theta} > \underline{c}(n)\} = \lim_{n \rightarrow \infty} n \mathbb{P}\{\bar{\theta} > \bar{c}(n)\} = 1.$$

From Lemma 5.2.2 in [14], we infer

$$0 < \lim_{n \rightarrow \infty} \frac{\mathbb{E}\underline{R}(n)^\beta}{\underline{c}^\beta(n)} < \infty \quad \text{and} \quad 0 < \lim_{n \rightarrow \infty} \frac{\mathbb{E}\bar{R}(n)^\beta}{\bar{c}^\beta(n)} < \infty,$$

and since

$$\mathbb{E}\underline{R}(n)^\beta \leq \mathbb{E}R(n)^\beta \leq \mathbb{E}\bar{R}(n)^\beta,$$

relation (34) follows if we can show that, for some $c_1, C_1 > 0$ and all $n \in \mathbb{N}$,

$$c_1 n^{-(1/\alpha+\varepsilon)\gamma(n^{1/\alpha+\varepsilon})/\alpha} \leq \frac{\underline{c}(n)}{n^{1/\alpha}} \quad \text{and} \quad \frac{\bar{c}(n)}{n^{1/\alpha}} \leq C_1 n^{(1/\alpha+\varepsilon)\gamma(n^{1/\alpha+\varepsilon})/\alpha}. \quad (36)$$

We prove only the first inequality in (34), for the second one follows in a similar manner. It is known that $(\underline{c}(n))$ is regularly varying with index $1/\alpha$. Hence, for large enough n , we

have $\underline{c}(n) \leq n^{1/\alpha+\varepsilon}$. On the other hand, using the monotonicity of $x \mapsto x^{\gamma(x)}$, we have

$$n \mathbb{P}\{\underline{\theta} > \underline{c}(n)\} \simeq n \underline{c}(n)^{-\alpha-\gamma(\underline{c}(n))} \geq (n^{-1/\alpha} \underline{c}(n))^{-\alpha} (n^{1/\alpha+\varepsilon})^{-\gamma(n^{1/\alpha+\varepsilon})},$$

and therefore

$$\limsup_{n \rightarrow \infty} (n^{-1/\alpha} \underline{c}(n))^{-\alpha} (n^{1/\alpha+\varepsilon})^{-\gamma(n^{1/\alpha+\varepsilon})} \leq 1,$$

yielding

$$c_1 n^{-(1/\alpha+\varepsilon)\gamma(n^{1/\alpha+\varepsilon})/\alpha} \leq \frac{\underline{c}(n)}{n^{1/\alpha}}$$

for some $c_1 > 0$.

To show (35), we argue in a similar manner. Set

$$\overline{M}(n) := \max_{1 \leq k \leq n} (-\overline{\theta}_k^{-1}), \quad \underline{M}(n) := \max_{1 \leq k \leq n} (-\underline{\theta}_k^{-1}), \quad n \in \mathbb{N}.$$

The Fisher-Tippett-Gnedenko theorem implies that $(-\overline{c}(n)\overline{M}(n))$ and $(-\underline{c}(n)\underline{M}(n))$ both converge weakly to a Weibull distribution. Moreover, Theorem 2.1 in [18] ensures that the moments of order β also converge, so

$$0 < \lim_{n \rightarrow \infty} \mathbb{E}\left(-\overline{c}(n)\overline{M}(n)\right)^\beta < \infty \quad \text{and} \quad 0 < \lim_{n \rightarrow \infty} \mathbb{E}\left(-\underline{c}(n)\underline{M}(n)\right)^\beta < \infty.$$

On the other hand,

$$\begin{aligned} \mathbb{E}\left(-\underline{c}(n)\underline{M}(n)\right)^\beta &= \mathbb{E}\left(\underline{c}(n) \min_{1 \leq k \leq n} \underline{\theta}_k^{-1}\right)^\beta \\ &= \mathbb{E}\left(\frac{\underline{c}(n)}{\max_{1 \leq k \leq n} \underline{\theta}_k}\right)^\beta \geq \mathbb{E}\left(\frac{\underline{c}(n)}{\max_{1 \leq k \leq n} \theta_k}\right)^\beta. \end{aligned}$$

and

$$\begin{aligned} \mathbb{E}\left(-\overline{c}(n)\overline{M}(n)\right)^\beta &= \mathbb{E}\left(\overline{c}(n) \min_{1 \leq k \leq n} \overline{\theta}_k^{-1}\right)^\beta \\ &= \mathbb{E}\left(\frac{\overline{c}(n)}{\max_{1 \leq k \leq n} \overline{\theta}_k}\right)^\beta \leq \mathbb{E}\left(\frac{\overline{c}(n)}{\max_{1 \leq k \leq n} \theta_k}\right)^\beta. \end{aligned}$$

Combining this with (36), we obtain (35). \square

The next result is the counterpart of Lemma 3.2 in the infinite-mean case.

Lemma 4.2 *Let $(\theta_k^{(n)})_{n \in \mathbb{N}, k \in \mathbb{N}}$ be an array of independent copies of a positive random variable θ which satisfies the assumptions of Lemma 4.1. For each $n \in \mathbb{N}$, define the increasing random walk*

$$R^{(n)}(0) := 0, \quad R^{(n)}(k) := \theta_1^{(n)} + \dots + \theta_k^{(n)}, \quad k \in \mathbb{N}.$$

Then, for arbitrary $T > 0$,

$$\sum_{n=1}^{\infty} \sup_{t \in [0, T]} \left| \alpha^n \log R^{(n)}(\lfloor e^{t\alpha^{-(n-1)}} \rfloor) - t \right| < \infty \quad a.s.$$

Proof. Put $m_n(t) := \lfloor e^{t\alpha^{-(n-1)}} \rfloor$ and $N_n := m_n(T)$. Then

$$\alpha^{n-1} \log m_n(t) \leq t < \alpha^{n-1} \log(m_n(t) + 1)$$

and therefore

$$\begin{aligned} & \sup_{t \in [0, T]} \left| \alpha^n \log R^{(n)}(\lfloor e^{t\alpha^{-(n-1)}} \rfloor) - t \right| \\ & \leq \sup_{t \in [0, T]} \left| \alpha^n \log R^{(n)}(m_n(t)) - \alpha^n \log(m_n(t))^{1/\alpha} \right| + \sup_{t \in [0, T]} \left| \alpha^{n-1} \log m_n(t) - t \right| \\ & \leq \alpha^n \sup_{1 \leq k \leq N_n} \left| \log \frac{R^{(n)}(k)}{k^{1/\alpha}} \right| + \alpha^{n-1} \log(1 + (m_n(t))^{-1}) \\ & \leq \alpha^n \sup_{1 \leq k \leq N_n} \left| \log \frac{R^{(n)}(k)}{k^{1/\alpha}} \right| + \alpha^{n-1} \log 2. \end{aligned}$$

Hence, we must show

$$\sum_{n=1}^{\infty} \alpha^n \sup_{1 \leq k \leq N_n} \left| \log \frac{R^{(n)}(k)}{k^{1/\alpha}} \right| < \infty \quad a.s.$$

which amounts to checking the following two relations:

$$\sum_{n=1}^{\infty} \alpha^n \sup_{1 \leq k \leq N_n} \left(\log^+ \frac{R^{(n)}(k)}{k^{1/\alpha}} \right) < \infty \quad a.s. \quad (37)$$

and

$$\sum_{n=1}^{\infty} \alpha^n \sup_{1 \leq k \leq N_n} \left(\log^- \frac{R^{(n)}(k)}{k^{1/\alpha}} \right) < \infty \quad a.s. \quad (38)$$

Fixing $\beta \in (0, \alpha)$, we obtain

$$\begin{aligned} \beta \mathbb{E} \left(\sup_{1 \leq k \leq N_n} \log^+ \frac{R^{(n)}(k)}{k^{1/\alpha}} \right) &= \mathbb{E} \left(\log \sup_{1 \leq k \leq N_n} \frac{(R^{(n)}(k))^\beta}{k^{\beta/\alpha}} \right) \\ &\leq \mathbb{E} \left(\log \sum_{j=0}^{\lfloor \log N_n \rfloor} \sup_{k \in (e^{-(j+1)} N_n, e^{-j} N_n]} \frac{(R^{(n)}(k))^\beta}{k^{\beta/\alpha}} \right) \\ &\leq \log \sum_{j=0}^{\lfloor T\alpha^{-(n-1)} \rfloor} \mathbb{E} \left(\sup_{k \in (e^{-(j+1)} N_n, e^{-j} N_n]} \frac{(R^{(n)}(k))^\beta}{k^{\beta/\alpha}} \right) \end{aligned}$$

where the last line follows from Jensen's inequality. For any $\varepsilon > 0$, we further infer with the help of Lemma 4.1 and (34)

$$\begin{aligned} \mathbb{E} \left(\sup_{k \in (e^{-(j+1)}N_n, e^{-j}N_n]} \frac{R^{(n)}(k)^\beta}{k^{\beta/\alpha}} \right) &\leq C \mathbb{E} \left(\frac{R^{(n)}(\lceil e^{-j}N_n \rceil)}{\lceil e^{-j}N_n \rceil^{1/\alpha}} \right)^\beta \\ &\leq C \left(\lceil e^{-j}N_n \rceil^{(1/\alpha+\varepsilon)\gamma(\lceil e^{-j}N_n \rceil^{1/\alpha+\varepsilon})} \right)^{\beta/\alpha}, \end{aligned}$$

where $C \in (0, \infty)$ denotes a suitable constant which here and hereafter may differ from line to line. By combining the previous estimates and using the monotonicity of $x \mapsto x^{\gamma(x)}$, we obtain

$$\begin{aligned} \beta \mathbb{E} \left(\sup_{1 \leq k \leq N_n} \log \frac{R^{(n)}(k)}{k^{1/\alpha}} \right) &\leq C + \log \sum_{j=0}^{\lceil T\alpha^{-(n-1)} \rceil} \left(\lceil e^{-j}N_n \rceil^{(1/\alpha+\varepsilon)\gamma(\lceil e^{-j}N_n \rceil^{1/\alpha+\varepsilon})} \right)^{\beta/\alpha} \\ &\leq C + \log \left(\left(\lceil T\alpha^{-(n-1)} \rceil + 1 \right) \left(N_n^{(1/\alpha+\varepsilon)\gamma(N_n^{1/\alpha+\varepsilon})} \right)^{\beta/\alpha} \right). \end{aligned}$$

Consequently, (37) follows from the inequality

$$\sum_{n=1}^{\infty} \alpha^n \log \left(N_n^{(1/\alpha+\varepsilon)\gamma(N_n^{1/\alpha+\varepsilon})} \right) \leq C \sum_{n=1}^{\infty} \gamma(N_n^{1/\alpha+\varepsilon})$$

and the fact that $\int_{x_0}^{\infty} \gamma(\exp(e^x)) dx < \infty$ implies (see calculations on p. 473 in [8])

$$\sum_{n=1}^{\infty} \gamma(e^{u\alpha^{-n}}) < \infty.$$

for any $u > 0$.

Equation (38) is verified along similar lines. Using $\log^- x = \log(x^{-1} \wedge 1) \leq \log(1+x^{-1})$, we obtain for arbitrary $\beta \in (0, \alpha)$

$$\begin{aligned} \beta \mathbb{E} \left(\sup_{1 \leq k \leq N_n} \log^- \frac{R^{(n)}(k)}{k^{1/\alpha}} \right) &\leq \beta \mathbb{E} \left(\log \left(1 + \sup_{1 \leq k \leq N_n} \left(\frac{k^{1/\alpha}}{R^{(n)}(k)} \right) \right) \right) \\ &\leq \mathbb{E} \log \left(1 + \sum_{j=0}^{\lfloor \log N_n \rfloor} \sup_{k \in (e^{-(j+1)}N_n, e^{-j}N_n]} \left(\frac{k^{\beta/\alpha}}{R^{(n)}(k)^\beta} \right) \right) \\ &\leq \log \left(1 + \sum_{j=0}^{\lfloor \log N_n \rfloor} \mathbb{E} \left(\sup_{k \in (e^{-(j+1)}N_n, e^{-j}N_n]} \left(\frac{k^{\beta/\alpha}}{R^{(n)}(k)^\beta} \right) \right) \right) \\ &\leq \log \left(1 + \sum_{j=0}^{\lfloor \log N_n \rfloor} \mathbb{E} \left(\frac{\lceil e^{-j}N_n \rceil^{\beta/\alpha}}{(R^{(n)}(\lceil e^{-(j+1)}N_n \rceil))^\beta} \right) \right) \\ &\leq \log \left(1 + \sum_{j=0}^{\lfloor \log N_n \rfloor} \mathbb{E} \left(\frac{\lceil e^{-j}N_n \rceil^{1/\alpha}}{\max_{k=1, \dots, \lceil e^{-(j+1)}N_n \rceil} \theta_k^{(n)}} \right)^\beta \right). \end{aligned}$$

By (35) in Lemma 4.1,

$$\mathbb{E} \left(\frac{\lceil e^{-j} N_n \rceil^{1/\alpha}}{\max_{k=1, \dots, \lceil e^{-(j+1)} N_n \rceil} \theta_k^{(n)}} \right)^\beta \leq C \left(\lceil e^{-(j+1)} N_n \rceil^{(1/\alpha + \varepsilon)\gamma} \lceil e^{-(j+1)} N_n \rceil^{1/\alpha + \varepsilon} \right)^{\beta/\alpha},$$

and this implies

$$\sum_{n=1}^{\infty} \alpha^n \mathbb{E} \sup_{1 \leq k \leq N_n} \left(\log^- \frac{R^{(n)}(k)}{k^{1/\alpha}} \right) < \infty$$

by the same argument as above. The proof of Lemma 4.2 is complete. \square

The counterpart of Proposition 3.3 is next.

Proposition 4.3 *Let $(R^{(n)}(k))_{k \in \mathbb{N}_0, n \in \mathbb{N}}$ be as in Lemma 4.2 and put*

$$h_n(t) := \alpha^n \log R^{(n)}(\lfloor e^{t\alpha^{-(n-1)}} \rfloor)$$

for $n \in \mathbb{N}$ and $t \geq 0$.

(B1) *There exists a D -valued random process $(Z_\infty^*(t))_{t \geq 0}$ such that*

$$(h_n \circ \dots \circ h_1(t))_{t \geq 0} \xrightarrow{n \rightarrow \infty} (Z_\infty^*(t))_{t \geq 0} \quad a.s.$$

in the space D endowed with the J_1 -topology. The process $(Z_\infty^(t))_{t \geq 0}$ is the a.s. limit of $(\alpha^n \log Z_n(t))_{t \geq 0}$ as $n \rightarrow \infty$, where $Z_n(t)$ denotes the number of descendants in generation n of ancestors $1, \dots, \lfloor e^t \rfloor$ in a GWP with generic offspring variable θ and countably many ancestors $1, 2, \dots$ in generation 0. The process $(Z_\infty^*(t))_{t \geq 0}$ has the following representation:*

$$Z_\infty^*(t) := Z_\infty^{(*,1)} \vee Z_\infty^{(*,2)} \vee \dots \vee Z_\infty^{(*, \lfloor e^t \rfloor)}, \quad t \geq 0,$$

with $(Z_\infty^{(,j)})_{j \in \mathbb{N}}$ denoting independent copies of Z_∞^* , the limit in (12).*

(B2) *For every fixed $k \in \mathbb{N}_0$ there exists a copy $(Z_{k,\infty}^*(t))_{t \geq 0}$ of the process $(Z_\infty^*(t))_{t \geq 0}$ such that*

$$(h_n \circ \dots \circ h_{k+1}(t))_{t \geq 0} \xrightarrow{n \rightarrow \infty} (\alpha^k Z_{k,\infty}^*(t\alpha^{-k}))_{t \geq 0} \quad a.s.$$

in the space D endowed with the J_1 -topology.

(B3) *As $t \rightarrow \infty$,*

$$\inf_{k \in \mathbb{N}} h_k \circ \dots \circ h_1(t) \rightarrow \infty \quad a.s.$$

(B4) *For any $T > 0$,*

$$\sup_{k \in \mathbb{N}} \sup_{t \in [0, T]} \left| h_{n+k} \circ \dots \circ h_{n+1}(t) - t \right| \xrightarrow{n \rightarrow \infty} 0 \quad a.s.$$

(B5) *The set $\{\alpha^k Z_{k,\infty}^*(t\alpha^{-k}) : t \geq 0, k \in \mathbb{N}\}$ is almost surely dense in $[0, \infty)$.*

Proof. (B1) The statement is equivalent to

$$\left(\alpha^n \log R^{(n)} \circ \dots \circ R^{(1)}(j) \right)_{j \in \mathbb{N}} \xrightarrow{n \rightarrow \infty} (Z_\infty^*(j))_{j \in \mathbb{N}} \quad a.s.$$

Introducing

$$\begin{aligned} Z_{n,1} &:= R^{(n)} \circ \dots \circ R^{(1)}(1), \\ Z_{n,j} &:= R^{(n)} \circ \dots \circ R^{(1)}(j) - R^{(n)} \circ \dots \circ R^{(1)}(j-1), \quad j = 2, 3, \dots, \end{aligned}$$

we obtain independent GWP's $(Z_{n,1})_{n \in \mathbb{N}}, (Z_{n,2})_{n \in \mathbb{N}}, \dots$ with generic offspring variable θ . By (12), the random variables

$$Z_{\infty}^{(*,j)} := \lim_{n \rightarrow \infty} \alpha^n \log Z_{n,j}, \quad j \in \mathbb{N}$$

exist a.s. in $(0, \infty)$ and are independent with the same distribution as Z_{∞}^* . It follows that

$$\begin{aligned} \alpha^n \log R^{(n)} \circ \dots \circ R^{(1)}(j) &= \alpha^n \log(Z_{n,1} + \dots + Z_{n,j}) \\ &\xrightarrow{n \rightarrow \infty} Z_{\infty}^{(*,1)} \vee \dots \vee Z_{\infty}^{(*,j)}, \end{aligned}$$

for any $j \in \mathbb{N}$ which completes the proof of (B1).

(B2) This is an immediate consequence of (B1) and the identity

$$h_n \circ \dots \circ h_{k+1}(t) = \alpha^k \alpha^{(n-k)} R^{(n)} \circ \dots \circ R^{(k+1)}(\lceil e^{t\alpha^{-k}} \rceil), \quad t \geq 0$$

valid for $n > k$.

(B3) Keeping the notation from (B1), we have

$$h_k \circ \dots \circ h_1(t) = \alpha^k \log \left(\sum_{j=1}^{\lfloor e^t \rfloor} Z_{k,j} \right), \quad t \geq 0,$$

and hence for all $t \geq 0$ and arbitrary $n_0 \in \mathbb{N}$

$$\begin{aligned} \inf_{k \in \mathbb{N}} h_k \circ \dots \circ h_1(t) &\geq \inf_{k \in \mathbb{N}} \max_{1 \leq j \leq \lfloor e^t \rfloor} (\alpha^k \log Z_{k,j}) \\ &= \left(\inf_{k > n_0} \max_{1 \leq j \leq \lfloor e^t \rfloor} (\alpha^k \log Z_{k,j}) \right) \wedge \left(\inf_{k \leq n_0} \max_{1 \leq j \leq \lfloor e^t \rfloor} (\alpha^k \log Z_{k,j}) \right) \\ &\geq \left(\max_{1 \leq j \leq \lfloor e^t \rfloor} \inf_{k > n_0} (\alpha^k \log Z_{k,j}) \right) \wedge \left(\max_{1 \leq j \leq \lfloor e^t \rfloor} \inf_{k \leq n_0} (\alpha^k \log Z_{k,j}) \right) \end{aligned}$$

by the minimax inequality. Since the $\inf_{k \leq n_0} (\alpha^k \log Z_{k,j}), j \in \mathbb{N}$, are iid with a law having unbounded support², i.e.

$$\mathbb{P} \left\{ \inf_{k \leq n_0} (\alpha^k \log Z_{k,1}) > z \right\} > 0$$

for every $z > 0$, we deduce that

$$\max_{1 \leq j \leq \lfloor e^t \rfloor} \inf_{k \leq n_0} (\alpha^k \log Z_{k,j}) \xrightarrow{t \rightarrow \infty} \rightarrow \infty \quad \text{a.s.}$$

² this is an easy consequence of the fact that θ has unbounded support in view of $\mathbb{E}\theta = \infty$

By the same arguments,

$$\max_{1 \leq j \leq \lfloor e^t \rfloor} \inf_{k > n_0} (\alpha^k \log Z_{k,j}) \xrightarrow{t \rightarrow \infty} \rightarrow \infty \quad \text{a.s.}$$

if we can show the existence of $n_0 \in \mathbb{N}$ such that

$$\mathbb{P}\left\{ \inf_{k \geq n_0} (\alpha^k \log Z_{k,1}) > z \right\} > 0 \quad (39)$$

for all $z > 0$. To this end, fix $z > 0$ and note that, by [8, Theorem 2], we have $a := \mathbb{P}\{Z_\infty^* > z + 1\} < 1$. On the other hand, we know from (B1) that

$$\lim_{n \rightarrow \infty} \mathbb{P}\left\{ \sup_{k \geq n} |\alpha^k \log Z_{k,1} - Z_\infty^*| \geq 1 \right\} = 0.$$

In particular, for each $0 < \delta < a$, we find $n_0 \in \mathbb{N}$ such that

$$\mathbb{P}\left\{ \sup_{k \geq n_0} |\alpha^k \log Z_{k,1} - Z_\infty^*| \geq 1 \right\} < \delta,$$

and so

$$\begin{aligned} \mathbb{P}\left\{ \inf_{k \geq n_0} (\alpha^k \log Z_{k,1}) \leq z \right\} &= \mathbb{P}\left\{ \inf_{k \geq n_0} (\alpha^k \log Z_{k,1}) \leq z, Z_\infty^* \leq z + 1 \right\} \\ &\quad + \mathbb{P}\left\{ \inf_{k \geq n_0} (\alpha^k \log Z_{k,1}) \leq z, Z_\infty^* > z + 1 \right\} \\ &\leq 1 - a + \mathbb{P}\left\{ \sup_{k \geq n_0} |\alpha^k \log Z_{k,1} - Z_\infty^*| \geq 1 \right\} \\ &\leq 1 - a + \delta < 1, \end{aligned}$$

which proves (39).

The proofs of (B4) and (B5) are omitted because they follow verbatim those of parts (A4) and (A5) of Proposition 3.3 (with Lemma 4.2 for part (B4) instead of Lemma 3.2 for part (A4) in Proposition 3.3). \square

4.2 Proof of Theorems 2.8, 2.9, 2.11 and 2.13, and Proposition 2.12

Define the random map $\phi : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ by

$$\phi((x_1, x_2, \dots)) := \alpha(x_{R(1)}, x_{R(2)}, \dots) \quad (40)$$

and further $\phi_n : \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}^{\mathbb{N}}$ for $n \in \mathbb{N}$ by

$$\phi_n((x_1, x_2, \dots)) := \alpha(x_{R^{(n)}(1)}, x_{R^{(n)}(2)}, \dots), \quad n \in \mathbb{N} \quad (41)$$

which are independent copies of ϕ (compare (24) and (25) in the finite-mean case).

Proof (of Theorem 2.8). The convergence in (13) is a consequence of (4) and part (B1) of Proposition 4.3. The SFPE (14) follows from the continuity of the map ϕ with respect to the product topology on $\mathbb{R}^{\mathbb{N}}$. \square

Proof (of Theorem 2.9). Note first that $(N_{\lfloor \exp(x\alpha^{-n}) \rfloor})_{x \in \mathbb{R}^+}$ is the counting process associated with the increasing random sequence $(\alpha^n \log S_k^{(n)})_{k \geq 1}$. By Theorem 2.8, the finite-dimensional distributions of this sequence converge as $n \rightarrow \infty$ to those of the non-decreasing random sequence $(Z_\infty^{(*,1)} \vee \dots \vee Z_\infty^{(*,k)})_{k \geq 1}$ with associated counting process $(N''(x))_{x \in \mathbb{R}^+}$. Also, for every fixed $n \in \mathbb{N}$,

$$\lim_{k \rightarrow \infty} \alpha^n \log S_k^{(n)} = \lim_{k \rightarrow \infty} (Z_\infty^{(*,1)} \vee \dots \vee Z_\infty^{(*,k)}) = \infty \quad \text{a.s.}$$

because $\xi > 0$ a.s. and Z_∞^* is not bounded from above. The latter statement follows from the SFPE $\alpha(Z_\infty^{(*,1)} \vee \dots \vee Z_\infty^{(*,\xi)}) \stackrel{d}{=} Z_\infty^*$. By Lemma 5.2 from the Appendix, this implies the weak convergence of the corresponding counting processes on D endowed with the M_1 -topology.

Proof (of Theorem 2.11). Here the assertion is implied by

$$\begin{aligned} \mathbb{P} \left\{ T(\lfloor e^{\alpha^{-n}x} \rfloor) - n \leq k \right\} &= \mathbb{P} \left\{ S_1^{(n+k)} > \lfloor e^{\alpha^{-n}x} \rfloor \right\} \\ &= \mathbb{P} \left\{ \alpha^{n+k} \log S_1^{(n+k)} > \alpha^k x \right\} \xrightarrow{n \rightarrow \infty} \mathbb{P} \{ Z_\infty^* > \alpha^k x \}. \quad \square \end{aligned}$$

Proof (of Proposition 2.12). If ξ has a Sibuya-distribution with parameter α , then the size Z_n of the associated GWP in generation n has a Sibuya distribution with parameter α^n because $f_\alpha \circ f_\beta = f_{\alpha\beta}$ and hence,

$$\mathbb{E}t^{Z_n} = f_\alpha \circ \dots \circ f_\alpha(t) = f_{\alpha^n}(t).$$

By Lemma 5.3 in Appendix, the random variable Z_∞^* from (12) has a standard exponential law. It remains to argue that

$$\sum_{j=1}^{\infty} \delta_{Z_\infty^{(*,1)} \vee \dots \vee Z_\infty^{(*,j)}} \stackrel{d}{=} \sum_{i=1}^{\infty} G_i \delta_{P_i}.$$

But a well-known consequence of the memoryless property of the exponential distribution is that the record values (taken without repetitions and denoted by P_1, P_2, \dots) in the sequence $(Z_\infty^{(*,j)})_{j \geq 1}$ form a Poisson point process (Tata's representation, [17, p. 69]). Given the record values P_1, P_2, \dots , the interrecord times G_1, G_2, \dots are independent and have geometric distributions with parameters $e^{-P_1}, e^{-P_2}, \dots$, respectively, by [17, Theorem 17.1, p. 77]. \square

Proof (of Theorem 2.13). Again our arguments follow along similar lines as those in the proof of Theorem 2.6 for the finite-mean case. Let $(X_1^{(0)}, X_2^{(0)}, \dots)$ be a solution to (14), i.e.

$$(X_1^{(0)}, X_2^{(0)}, \dots) \stackrel{d}{=} \alpha(X_{R(1)}^{(0)}, X_{R(2)}^{(0)}, \dots).$$

By the Kolmogorov consistency theorem, the underlying probability space $(\Omega, \mathfrak{A}, \mathbb{P})$ may be assumed to be large enough to carry the following objects:

- the random sequence $(X_1^{(0)}, X_2^{(0)}, \dots)$;
- a two-sided sequence $(R^{(n)}(\cdot))_{n \in \mathbb{Z}}$ of independent copies of the random walk $R(\cdot)$ and the corresponding sequence of random maps (41);
- a two-sided stationary sequence $(V_k)_{k \in \mathbb{Z}} = ((X_1^{(k)}, X_2^{(k)}, \dots))_{k \in \mathbb{Z}}$ such that V_k is independent of $(\phi_n)_{n \leq k}$ for each $k \in \mathbb{Z}$, and

$$V_0 := (X_1^{(0)}, X_2^{(0)}, \dots) \quad \text{and} \quad V_k = \phi_{k+1}(V_{k+1}), \quad k \in \mathbb{Z},$$

thus

$$(X_1^{(k)}, X_2^{(k)}, \dots) = \alpha \left(X_{R^{(k+1)}(1)}^{(k+1)}, X_{R^{(k+1)}(2)}^{(k+1)}, \dots \right), \quad k \in \mathbb{Z}.$$

Define a sequence of random measures $(\nu_n)_{n \geq 0}$ on $[0, \infty)$ by

$$\nu_n[0, t] := \alpha^n \log X_{[\exp(t\alpha^{-n})]}^{(n)}, \quad t \geq 0$$

for $n \in \mathbb{N}_0$. As in the proof of Theorem 2.6, it is enough to show that the sequence $(\nu_n)_{n \in \mathbb{N}}$ converges almost surely in the vague topology, i.e.

$$\nu_n \xrightarrow{v} \nu_\infty \quad \text{a.s.} \quad (42)$$

Recall from Proposition 4.3 that $h_n(t) = \alpha^n \log R^{(n)}(\lfloor e^{t\alpha^{-(n-1)}} \rfloor)$ for $n \in \mathbb{N}$ and $t \geq 0$. We have

$$\nu_n[0, t] = \nu_{n+1}[0, h_{n+1}(t)] \quad (43)$$

and upon iteration

$$\nu_k[0, t] = \nu_{n+k}[0, h_{n+k} \circ \dots \circ h_{k+1}(t)] \quad (44)$$

for $n \in \mathbb{N}_0$ and $t \geq 0$, in particular for $k = 0$

$$\nu_0([0, t]) = \nu_n \left[0, \alpha^n \log \left(\sum_{j=1}^{\lfloor e^t \rfloor} Z_{n,j} \right) \right], \quad n \in \mathbb{N}, \quad t \geq 0. \quad (45)$$

Since

$$\lim_{t \rightarrow \infty} \lim_{n \rightarrow \infty} \alpha^n \log \sum_{j=1}^{\lfloor e^t \rfloor} Z_{n,j} = \lim_{t \rightarrow \infty} Z_\infty^*(t) = \infty \quad \text{a.s.},$$

see (B1) in Proposition 4.3, equation (45) implies that, for any $T > 0$,

$$\sup_{n \in \mathbb{N}_0} \nu_n[0, T] < \infty \quad \text{a.s.}$$

and therefore almost sure relative compactness of $(\nu_n)_{n \geq 0}$ in the vague topology.

Let $(\nu_{m_n})_{n \geq 1}$ be an a.s. vaguely convergent subsequence and ν'_∞ its limit. From (44), we have a.s. for each $k \in \mathbb{N}_0$ and $m_n > k$ that

$$v_k[0, t] = v_{m_n}[0, h_{m_n} \circ \dots \circ h_{k+1}(t)], \quad t \geq 0. \quad (46)$$

By part (B2) of Proposition 4.3,

$$h_{m_n} \circ \dots \circ h_{k+1}(t) \xrightarrow{n \rightarrow \infty} \alpha^k Z_{k, \infty}^*(t\alpha^{-k}) \quad \text{a.s.}$$

in the space D endowed with the J_1 -topology. Sending $n \rightarrow \infty$ in (46) and applying Lemma 5.1, we obtain

$$v_k[0, t] = v'_\infty[0, \alpha^k Z_{k, \infty}^*(t\alpha^{-k})], \quad t > 0$$

for every fixed $k \in \mathbb{N}_0$. By part (B6) of Proposition 4.3, the random set consisting of $\alpha^k Z_{k, \infty}^*(t\alpha^{-k})$ for $t \geq 0$, $k \in \mathbb{N}_0$ is a.s. dense in $[0, \infty)$. Hence, if v''_∞ denotes another subsequential limit of $(v_n)_{n \geq 1}$, then a.s.

$$v''_\infty[0, t] = v'_\infty[0, t]$$

on a dense subset of $[0, \infty)$ and therefore $v''_\infty \equiv v'_\infty$, proving the convergence (42).

The restricted self-similarity property (15) can be checked as in the proof of Theorem 2.6. \square

5 Appendix

Lemma 5.1 *Let $\mathcal{M} := \mathcal{M}[0, \infty)$ be the set of locally finite measures on $[0, \infty)$ endowed with the vague topology and let $\phi : \mathcal{M} \times [0, \infty) \rightarrow \mathbb{R}^+$ be defined by*

$$\phi(\mu, x) = \mu([0, x]).$$

Endowing $\mathcal{M} \times [0, \infty)$ with the product topology, the mapping ϕ is continuous at any (μ, x) such that $\mu(\{x\}) = 0$.

Proof. Let $\mu_n \xrightarrow{v} \mu_0$, $\lim_{n \rightarrow \infty} x_n = x_0$ and set $f_0(s) := \mathbf{1}_{\{s \leq x_0\}}$, $B_0 := (0, \infty) \setminus \{x_0\}$, $f_n(s) := \mathbf{1}_{\{s \leq x_n\}}$, $B_n := [0, \infty)$ for $n \in \mathbb{N}$. The result now follows from Lemma 15.7.3 in [16]. \square

Lemma 5.2 *For every $n \in \mathbb{N} \cup \{\infty\}$ let $0 \leq X_1^{(n)} \leq X_2^{(n)} \leq \dots$ be a random sequence such that $\lim_{k \rightarrow \infty} X_k^{(n)} = \infty$ a.s. and*

$$(X_1^{(n)}, X_2^{(n)}, \dots) \xrightarrow{f.d.d.} (X_1^{(\infty)}, X_2^{(\infty)}, \dots). \quad (47)$$

Define the corresponding counting processes $N^{(n)}(x) := \#\{k \in \mathbb{N} : X_k^{(n)} \leq x\}$, $x \geq 0$, $n \in \mathbb{N} \cup \{\infty\}$. Then

$$(N^{(n)}(x))_{x \geq 0} \xrightarrow{d} (N^{(\infty)}(x))_{x \geq 0}$$

in D endowed with the M_1 -topology, and the latter may be even replaced with the J_1 -topology if the sequence $(X_k^{(\infty)})_{k \geq 1}$ is strictly increasing with probability 1.

Proof. Consider the space L_{\leq} of all sequences $y = (y_k)_{k \in \mathbb{N}}$ such that $0 \leq y_1 \leq y_2 \leq \dots$ and $\lim_{k \rightarrow \infty} y_k = \infty$. We endow L_{\leq} with the topology of pointwise convergence. It is well-known that this topology is metrizable. Consider a map $\Psi : L_{\leq} \rightarrow D$ which assigns to each sequence $y = (y_k)_{k \in \mathbb{N}}$ the corresponding counting function

$$\Psi(y)(x) = \#\{k \in \mathbb{N} : y_k \leq x\}, \quad x \geq 0.$$

If we endow the D with the M_1 -topology, then it is an easy exercise to check that the map Ψ is continuous on L_{\leq} . Let us now endow D with the J_1 -topology. Then, Ψ is not continuous on L_{\leq} (because the points can build clusters in the limit), but it is continuous on the subset $L_{<}$ consisting of strictly increasing sequences.

We can consider $(X_k^{(n)})_{k \geq 1}$, $n \in \mathbb{N} \cup \{\infty\}$, as random elements with values in L_{\leq} (in $L_{<}$ when using the J_1 -topology). Then the convergence in (47) is equivalent to the weak convergence of the corresponding random elements. The statement of the lemma thus follows from the continuous mapping theorem. \square

Lemma 5.3 *If S_α denotes a random variable with a Sibuya distribution with parameter $0 < \alpha < 1$, then*

$$\lim_{\alpha \downarrow 0} \mathbb{P}\{\alpha \log S_\alpha \leq x\} = 1 - e^{-x}$$

for all $x > 0$.

Remark 5.4 *It is known [6] that if Z_α is a positive α -stable variable with $\alpha \in (0, 1)$, then $\alpha \log Z_\alpha$ converges in distribution to the Gumbel double exponential law.*

Proof (Proof of Lemma 5.3). Consider independent Bernoulli variables B_1, B_2, \dots , such that $\mathbb{P}\{B_i = 1\} = \alpha/i$. Then it is easy to check that $S_\alpha := \min\{n \in \mathbb{N} : B_i = 1\}$ has the required Sibuya distribution with parameter α . Taking any $x > 0$, we now infer

$$\mathbb{P}\{\alpha \log S_\alpha \leq x\} = \mathbb{P}\{S_\alpha \leq \lfloor e^{x/\alpha} \rfloor\} = 1 - \mathbb{P}\{B_1 + \dots + B_{\lfloor e^{x/\alpha} \rfloor} = 0\}.$$

But for $\alpha \downarrow 0$, the probability on the right-hand side converges to e^{-x} because

$$B_1 + \dots + B_{\lfloor e^{x/\alpha} \rfloor} \xrightarrow[\alpha \downarrow 0]{d} \text{Poi}(x).$$

For the proof, just note that $\sum_{i=1}^{\lfloor e^{x/\alpha} \rfloor} \frac{\alpha}{i} \simeq \alpha \log \lfloor e^{x/\alpha} \rfloor \simeq x$, as $\alpha \downarrow 0$, and apply the Poisson limit theorem. \square

Acknowledgements The research of Gerold Alsmeyer was supported by the Deutsche Forschungsgemeinschaft (SFB 878), the research of Alexander Marynych by the Alexander von Humboldt Foundation.

References

1. G. Alsmeyer, Z. Kabluchko, and A. Marynych. A leader-election procedure using records, 2015. Preprint available at <http://arxiv.org/abs/1512.04750>.

2. K. B. Athreya. Coalescence in the recent past in rapidly growing populations. *Stochastic Process. Appl.*, 122(11):3757–3766, 2012.
3. K. B. Athreya and P. E. Ney. *Branching processes*. Die Grundlehren der mathematischen Wissenschaften, Band 196. Springer-Verlag, New York, 1972.
4. N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
5. F. T. Bruss and R. Grübel. On the multiplicity of the maximum in a discrete random sample. *Ann. Appl. Probab.*, 13(4):1252–1263, 2003.
6. N. Cressie. A note on the behaviour of the stable distributions for small index α . *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 33(1):61–64, 1975/76.
7. D. A. Darling. The Galton-Watson process with infinite mean. *J. Appl. Probability*, 7:455–456, 1970.
8. P. L. Davies. The simple branching process: a note on convergence when the mean is infinite. *J. Appl. Probab.*, 15(3):466–480, 1978.
9. Y. Davydov, I. Molchanov, and S. Zuyev. Strictly stable distributions on convex cones. *Electron. J. Probab.*, 13:no. 11, 259–321, 2008.
10. Y. Davydov, I. Molchanov, and S. Zuyev. Stability for random measures, point processes and discrete semigroups. *Bernoulli*, 17(3):1015–1043, 2011.
11. J. A. Fill, H. M. Mahmoud, and W. Szpankowski. On the distribution for the duration of a randomized leader election algorithm. *Ann. Appl. Probab.*, 6(4):1260–1283, 1996.
12. D. R. Grey. Almost sure convergence in Markov branching processes with infinite mean. *J. Appl. Probability*, 14(4):702–716, 1977.
13. R. Grübel and K. Hagemann. Leader election: A Markov chain approach. *Mathematica Applicanda*, 44(1):113–134, 2016.
14. I. A. Ibragimov and Y. V. Linnik. *Independent and stationary sequences of random variables*. Wolters-Noordhoff Publishing, Groningen, 1971. With a supplementary chapter by I. A. Ibragimov and V. V. Petrov, Translation from the Russian edited by J. F. C. Kingman.
15. S. Janson and W. Szpankowski. Analysis of an asymmetric leader election algorithm. *Electron. J. Combin.*, 4(1):Research Paper 17, 16 pp. (electronic), 1997.
16. O. Kallenberg. *Random measures*. Akademie-Verlag, Berlin; Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London, 3rd edition, 1983.
17. V. B. Nevzorov. *Records: mathematical theory*, volume 194 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 2001. Translated from the Russian manuscript by D. M. Chibisov.
18. J. Pickands, III. Moment convergence of sample extremes. *Ann. Math. Statist.*, 39:881–889, 1968.
19. H. Prodinger. How to select a loser. *Discrete Math.*, 120(1-3):149–159, 1993.
20. H.-J. Schuh and A. D. Barbour. On the asymptotic behaviour of branching processes with infinite mean. *Advances in Appl. Probability*, 9(4):681–723, 1977.
21. E. Seneta. The simple branching process with infinite mean. I. *J. Appl. Probability*, 10:206–212, 1973.
22. R. van der Hofstad, G. Hooghiemstra, and D. Znamenski. Distances in random graphs with finite mean and infinite variance degrees. *Electron. J. Probab.*, 12:no. 25, 703–766, 2007.
23. B. von Bahr and C.-G. Esseen. Inequalities for the r th absolute moment of a sum of random variables, $1 \leq r \leq 2$. *Ann. Math. Statist.*, 36:299–303, 1965.
24. G. Zanella and S. Zuyev. Branching-stable point processes, 2016. Preprint available at <http://arxiv.org/pdf/1503.01329v2.pdf>.